

The effects of UV photometry and binary interactions on photometric redshift and galaxy morphology

Fenghui Zhang^{1*}, Zhanwen Han¹, Lifang Li¹, Hongguang Shan¹ and Yu Zhang^{1,2}

¹National Astronomical Observatories/Yunnan Observatory, Chinese Academy of Sciences, Kunming, 650011, China

²Graduate University of the Chinese Academy of Science, Beijing 100049, China

24 June 2010

ABSTRACT

Using the *Hyperz* code (Bolzonella et al. 2000) and a template spectral library which consists of 4 observed galaxy spectra from Coleman, Wu & Weedman (CWW, 1980) and 8 spectral families built with evolutionary population synthesis models, we present photometric redshift estimates (photo- z) for a spectroscopic sample of 6,531 galaxies, for which spectroscopic redshifts are also available and are selected from the SDSS DR7 and GALEX DR4, and morphologies for a morphological sample of 1,502 bright galaxies, which are from the catalogue of Fukugita et al. (2007) and also matched with the SDSS DR7 and GALEX DR4.

We find that the inclusion of F_{UV} or N_{UV} or both photometry decreases the number of catastrophic identifications ($|z_{\text{phot}} - z_{\text{spec}}| > 1.0$). If catastrophic identifications are removed, the inclusion of both F_{UV} and N_{UV} photometry mainly increases the number of non-catastrophic identifications in the low redshift, $g - r \lesssim 0.8$ and fainter r -magnitude regions. The inclusion of binary interactions mainly increases the number of non-catastrophic identifications and decreases the deviations in the $0.3 \leq g - r \leq 0.8$ region in the case of only using optical photometry.

Based on the morphological galaxy sample, we find that the inclusion of UV photometry would decrease and increase the probability that early types are classified as Burst and E types, respectively, and increase that late types are classified as CWW-Sbc and CWW-Scd types. If catastrophic identifications are excluded, the inclusion of UV data mainly raises the identifications of late types in all redshift, bluer $g - r$ and $r \gtrsim 14$ regions. Moreover, Binary interactions mainly affect the determinations of E and S0 types.

By comparison we find that the *reliability* and *completeness* for early- and late-type selection by the *Hyperz* code are less than those by the concentration index $C' = 2.6$, profile likelihood $P_{\text{exp}} - P_{\text{dev}} = 0$ and colour $u - r = 2.22$ criteria. Moreover, we find that $N_{UV} - u = 1.94$ and $5.77 - 1.47(u - r) = F_{UV} - u$ discriminators can be used as morphology selection indicators. These two criteria have comparable *reliability* and *completeness* for selecting early- and late-type galaxies to $C' = 2.6$ criterion and higher *completeness* for early-type selection than $u - r = 2.22$ criterion.

Key words: Galaxies: distances and redshifts – Galaxies: fundamental parameters – binary: general – ultraviolet: galaxies

1 INTRODUCTION

Redshift is one of the key ingredients of cosmology, it can be used to estimate distances and hence to place observed properties on a physical scale. Many modern astrophysical measurements and researches (identifications of galaxy clusters and very high redshift objects, baryon acoustic oscillation measurements, galaxy evolution, large scale structure

and gravitational lensing studies) have benefited from substantially larger catalogues of redshifts (Gerdes et al. 2009).

Galaxy redshift plays an important role in studies of galaxy formation and evolution. Up to now, the spectroscopic redshifts of many relatively bright galaxies are obtained, while those of fainter galaxies are difficult to obtain. Because that photometry is ~ 2 orders of magnitude lesser time-consuming than spectroscopy for a given telescope size, the redshifts of fainter galaxies ($R \gtrsim 25$ mag) or upcoming surveys will exclusively rely on the broad-band,

* E-mail: gssephd@public.km.yn.cn; zhang-fh@hotmail.com

medium-band or custom-designed narrow-band photometry (Brammer, van Dokkum & Coppi 2008; Niemack et al. 2009, and references therein).

The technique for deriving redshifts from broad-band photometry was pioneered by Baum (1962). So far, a number of codes have been developed to compute photometric redshifts (photo- z s), such as *BPZ* (Bentz 2000), *Hyperz* (Bolzonella, Miralles & Pelló 2000), *GREGZ*¹ (Rudnick et al. 2001, 2003), *IMPZ* (Baddedge et al. 2004), *LePHARE*² (Arnouts & Ilbert), *ZEBRA* (Feldmann et al. 2006), *kcorrect* (Blanton & Roweis 2007), *LRT* (Assef et al. 2008), *EAZY* (Brammer et al. 2008), *ARBORZ* (Gerdes et al. 2009), Artificial Neural Networks (ANNs) and so on. Three kinds of algorithm have been used in the photo- z calculations (Abdalla et al. 2008; Brammer et al. 2008; Gerdes et al. 2009; Niemack et al. 2009; Zhang et al. 2009).

- The template fitting approach has even been used. In this scheme photo- z is obtained by comparing the observed spectra with a set of template spectra, which can be based on either population synthesis models or empirical studies.

- The empirical training set approach has been used. The basic principle of this method is the derivation of a parameterization of redshift through the magnitudes of the galaxies in a training set, this parameterization is then applied to galaxies whose redshifts need to be estimated, yielding an estimation of the photometric redshift. The galaxies in the training set have known spectroscopic redshifts and similar properties to the galaxies for which one wants to estimate the redshifts. ANNs is an empirical training set approach.

- Instance-based learning method also has been used. This method also relies on the real data.

Among the traditional template fitting codes, some of them (such as *Hyperz*, *IMPZ*, *LePHARE*, etc) used a straightforward χ^2 minimization algorithm to obtain photo- z . As spectral information provided by broadband photometry is at low resolution, the photo- z s of galaxies have relatively large errors. To avoid catastrophic errors, some codes improved the template spectra, for example, Niemack et al. (2009) developed a set of high-resolution spectral templates based on galaxy physical information about the star-formation history, other codes used a refinement of the χ^2 minimization method, for example, (i) *BPZ* code incorporated Bayesian statistics and allowed the use of extra information of galaxies (priors), (ii) *EASY*, *ZEBRA* and *kcorrect* codes used the hybrid method: *EASY* code allowed the linear combination of templates as done in *GREGZ* code and the use of priors; *ZEBRA* code used the combination of the priors [which were used to calculate a prior self-consistently from the photometric catalogue when it is run in Bayesian mode] with the training set method [which was used to improve the standard set of templates in specified redshift bins in its template optimization mode by the advantage of the zCOSMOS database (Lilly et al. 2007)]; *kcorrect* code used the data of Sloan Digital Sky Survey (SDSS, York et al. 2000) project and incorporated the

principle component analysis method to optimize the template. Among these photo- z codes, the priors of luminosity function (Mobasher et al. 2007) and surface luminosity (Xia et al. 2009) have been used.

Among those non-traditional photometry fitting codes, some of them included the structural properties of galaxies, for example, Wray & Gunn (2008) used surface brightness and the Sérsic index (a measurement of the radial light profile) in addition to five-band photometry to obtain photo- z s, and Sarajedini et al. (1999) used bulge-to-total flux ratio along with I -magnitude and $V - I$ colour to obtain photo- z s. Negrello et al. (2009) incorporated the features of polycyclic aromatic hydrocarbons and silicate features into mid-infrared wavelength window to recover the photo- z s of starburst galaxies. Moreover, Kurtz et al. (2007, μ -photoZ) even only used one colour and the surface brightness from a single band to obtain photo- z s of galaxies.

Now, photo- z s have been extensively used in many studies, such as they were used in deep cosmological surveys to yield the galaxy luminosity function and the evolution of star formation rate (SFR) (Mobasher et al. 2007, and references therein).

Morphology is another important parameter of galaxy. First, it is one of the first segregation of galaxy clusters, different morphological types exhibit distinctly different astrophysical properties, reflecting the different histories of the formation and evolution of galaxies (Shimasaku et al. 2001). Studying it can help us to investigate the properties of classified galaxies [such as colours, the absence/presence of emission lines or the galaxy profiles (Lintott et al. 2008)]. Moreover, studying the correlation of morphology with other galaxy properties (such as luminosity function, Blanton et al. 2001) can help us to understand the physical processes in galaxies (such as the influence of star formation on the process of galaxy formation) and the theories of galaxy formation and evolution (Blanton et al. 2001). So far, several approaches and techniques have been proposed to determine the morphological types.

- Automated classification approach, i.e., using the parameter(s) sensitive to morphology to obtain automatically the galaxy morphology, has been used. This method spends less time, but faces on a serious problem that it is difficult to find such parameter(s). In history, the following parameters have been used in the automated classification method: $u - r$ colour (Strateva et al. 2001), profile probabilities (the relation between exponential P_{exp} and de Vaucouleurs' P_{dev} profile likelihoods), concentration index (the ratio of the radii containing 90% to 50% of the *Petrosian* r galaxy light, $C \equiv r_{p90}/r_{p50}$, Shimasaku et al. 2001; Strateva et al. 2001) and so on.

- Visual classification approach, i.e., classifying galaxies by visual inspection of experts or volunteers, has been used. This method serves as the most reliable method when adopting the Hubble classification (Sandage 1961) for galaxies with large apparent sizes (Shimasaku et al. 2001). The greatest disadvantage of this method is more time-consuming, the size of the galaxy sample is small and the galaxies inspected are relatively bright. Fukugita et al. (2007) have compiled a catalogue about morphological classification of ~ 2500 bright galaxies in the SDSS Data Re-

¹ Because Greg Rudnick did not name his code, we follow the nomenclature of Brammer et al. (2008)

² See <http://www.oamp.fr/people/arnouts/LEPHARE.html>.

lease three (DR3) by the visual inspection of three expert classifiers. Galaxy Zoo project has invited a large number of volunteers to separate early-type galaxies from spirals in the large data sets by using proxies for morphology (Lintott et al. 2008).

- Spectrum fitting approach, i.e., fitting the observed spectra to the template spectra of galaxies with known morphological types, has been used.
- The k -means method also has been used (Zhang et al. 2009).

We will use one of the widely adopted template-fitting codes, *Hyperz*, to obtain the photo- z s of galaxies. Meanwhile the rough morphological types will also be obtained because the *Hyperz* code allows to obtain the best set of fit parameters. As the first step of this work, we will use *Hyperz* code to obtain the photo- z s and morphological types of galaxies with known spectroscopic redshifts and morphological types. This can be used to explore the ability of the *Hyperz* code of recovering redshifts and morphological types of galaxies, and judge whether or how the recovered parameters are affected by the input parameters (including ultraviolet [UV] photometry, binary interactions and population synthesis models).

The reasons of discussing the influence of UV photometry and binary interactions on the determinations of photo- z s and morphological types are as follows.

(i) In those previous works using template fitting method to obtain photo- z s of galaxies, the template spectra usually came from real objects or population synthesis models of single stellar populations (SSPs), rather than the population synthesis models including binary interactions. As for population synthesis models of binary stellar populations (BSPs), the Yunnan group (Zhang et al. 2004b, 2005) have considered various binary interactions in evolutionary population synthesis (EPS) models, and drawn the conclusion that the inclusion of binary interactions can affect the overall shape of the spectral energy distribution (SED) of population, in particular, the SED of population in the UV passbands is bluer by 2-3 mag at $t \sim 1$ Gyr if binary interactions are accounted for. Above conclusion further has been confirmed by Han, Podsiadlowski & Lynas-Gray (2007, hereafter HPL07), who included hot subdwarf B stars (sdBs) in population. The bluer SED in the UV passbands can be explained by that binary interactions can create some important classes of objects, such as sdBs for a population older than ~ 1 Gyr (HPL07) and blue stragglers at 0.5–1.5 Gyr.

(ii) Observations find that almost all elliptical galaxies have UV-upturn phenomenon, i.e., the flux increases with the wavelength decreasing from 2000 to 1200 Å (HPL07). This phenomenon can be explained by the EPS models of BSPs (HPL07). Thus, if only optical photometry and/or the EPS models of SSPs are used to obtain photo- z s and morphologies of elliptical galaxies, whether these results are reasonable or not?

(iii) Another fact is that the photo- z s, which are derived by using the *Hyperz* code under the condition of neglecting UV photometry, significantly deviate from the spectroscopic ones for some galaxies in the redshift range $z < 0.2$ (also see Fig. 4).




Figure 1. Comparison of SED among BC03 (red dotted line), Yunnan (black solid line + triangles for Yunnan-b; black solid line for Yunnan-s) and HPL07 models (green dot-dashed line + circles for HPL07-b; green dot-dashed line for HPL07-s) for a solar-metallicity population (normalized to $1 M_{\odot}$) at $\log(t) = 8.85$ yr.

In this work, we use two samples of galaxies: spectroscopic and morphological. The galaxies in the spectroscopic sample are selected randomly from the SDSS DR7 and their spectroscopic redshifts are available. The galaxies in the morphological sample are from the catalogue of Fukugita et al. (2007), which presented the morphological types of 2,253 bright galaxies in the SDSS DR3 by using independent classification scheme (visual classification). All galaxies in both samples have matched with the Galaxy Evolution Explorer (GALEX) DR4 (see Section 3 for details). Moreover, we not only use the population synthesis models of SSPs but also those of BSPs to construct theoretical template library (see Section 2). Niemack et al. (2009) have used the SDSS and GALEX photometric data to study the redshifts of galaxies, but they used the population synthesis models of SSPs.

The outline of the paper is as follows. In Sections 2 and 3 we describe the method used to obtain photo- z and the galaxy samples, respectively. In Sections 4 and 5 we obtain photo- z estimates for the spectroscopic galaxy sample and morphologies for the morphological galaxy sample, and discuss the effects of UV photometry and binary interactions on them. Finally we present a summary and conclusions in Section 6.

2 METHOD

The photo- z and galaxy morphology are computed through the *Hyperz* code of Bolzonella et al. (2000), which is the first publicly available photo- z code and has been widely used in the literature for photo- z estimations of galaxies (Abdalla et al. 2008). *Hyperz* is a template fitting procedure and adopts a standard χ^2 minimization algorithm:

Table 1. Description about GISSSEL98, BC03, HPL07 and Yunnan spectral synthesis models.

population	GISSSEL98&BC03 (SSP)	HPL07 (SSP&BSP)	Yunnan (SSP&BSP)
Ages	221	88	90
$\log(t_i/\text{yr})$	5.100	8.000	5.000
$\log(t_f/\text{yr})$	10.300	10.175	10.175
$\Delta\log(t/\text{yr})$	0.050 [5.100 ≤ $\log(t/\text{yr})$ ≤ 6.000; 7.757 ≤ $\log(t/\text{yr})$ ≤ 9.207] 0.020 [6.000 ≤ $\log(t/\text{yr})$ ≤ 7.440] 0.005~0.038 [7.440 ≤ $\log(t/\text{yr})$ ≤ 7.757; $\log(t/\text{yr})$ ≥ 9.207]	0.025	0.100 [5.000 ≤ $\log(t/\text{yr})$ ≤ 6.500] 0.050 [$\log(t/\text{yr})$ ≥ 6.500]

$$\chi^2 = \sum_{i=1}^{N_{\text{filters}}} \left[\frac{F_{\text{obs},i} - b \times F_{\text{temp},i}}{\sigma_i} \right]^2 \quad (1)$$

where $F_{\text{obs},i}$, $F_{\text{temp},i}$ and σ_i are the observed and template fluxes and their uncertainty in filter i , respectively, and b is a normalization constant.

Hyperz code is based on the fit of the overall shape of the spectra and on the detection of strong spectral features, such as the 4000 Å break, Balmer break, Lyman decrement or strong emission lines, and adopts a standard SED fitting method.

Hyperz takes as its inputs the filter set and the photometric catalogue of galaxies which comprises magnitudes and photometric errors through the filters specified in the filter set. For a given filter set and galaxy catalogue, the relevant parameters introduced in the photo- z calculation are: (i) the set of template spectra, which can be observed SEDs or built with spectral models or both (see this section), if spectral models are used, the type of SFR, the possible link between the age and the metallicity of the stellar population, and the choice of an initial mass function (IMF) are involved; (ii) the reddening law (*Hyperz* code provides 5 laws to choose); (iii) flux decrements in the Lyman forest; (iv) the limiting magnitude in each filter; and, (v) the cosmological parameters H_0 , Ω_M and Ω_Λ , and so on.

The descriptions about the filter set and the photometric catalogue of galaxies are presented in Section 3. This section describes emphatically the template spectral library (Sections 2.1 and 2.2) and the definition of models (see Section 2.3).

2.1 Spectral synthesis models

In the *Hyperz* package observed SEDs and the spectral synthesis models of GISSSEL98 (Galaxy Isochrone Synthesis Spectral Evolution Library, Bruzual & Charlot 1993) are provided. To check the effects of spectral synthesis models and binary interactions on the determinations of photo- z s and morphologies of galaxies, we also build the theoretical template SEDs by using the Bruzual & Charlot (2003, hereafter BC03), HPL07 and Yunnan models (Zhang et al. 2002, 2004a,b, 2005).

GISSSEL98 and BC03 models were built by Bruzual & Charlot in 1993 and 2003, respectively. In this work the GISSSEL98 models with the IMF by Miller & Scalo (1979) and the BC03 models, which use the Padova 1994 stellar evolutionary tracks and the Chabrier (2003) IMF, are used because the choice of IMF has a negligible impact on the

final results (Bolzonella et al. 2000). Both models did not take binary interactions into account.

HPL07 models included the binary interactions of sdBs (Han et al. 2002, 2003) in populations. The Yunnan models were developed by Zhang and her colleagues, in their models the rapid single/binary evolution codes (Hurley, Pols & Tout 2000; Hurley, Tout & Pols 2002) were used for the single/binary evolutionary tracks (the rapid binary evolution code includes those binary interactions studied before 2002). Both models presented the SEDs of populations without and with binary interactions, i.e., SSPs and BSPs. To distinguish them we name the HPL07 models for SSPs and BSPs HPL07-s and HPL07-b, name the Yunnan models for SSPs and BSPs Yunnan-s and Yunnan-b models, respectively. Detailed description of IMFs used by HPL07 and Yunnan models has presented in their related papers, we refer the interested reader to them.

Moreover, the ages and age interval of these EPS models are different. In Table 1 we give the initial and final ages [$\log(t_i/\text{yr})$, $\log(t_f/\text{yr})$], the age interval [$\Delta\log(t/\text{yr})$] within a given age range (in the brackets) for these EPS models. In Fig. 1 we give a comparison in the SED of a solar-metallicity population (normalized to $1M_\odot$) among BC03, HPL07 and Yunnan models at $\log(t/\text{yr}) = 8.85$. From it we see that the UV spectra of SSP differs significantly from that of BSP for both HPL07 and Yunnan models, and the difference in the SED among BC03, HPL07-s and Yunnan-s models is also large. Therefore, it is necessary to discuss the effects of spectral synthesis models, UV photometry and binary interactions on the determinations of photo- z s and morphological types of galaxies.

2.2 Theoretical template SED library

The template spectral library should comprise the SEDs with different ages and spectral types of galaxies. Spectral synthesis models only provide the SEDs of populations without any SFR at different ages. Therefore, at a given age we need to generate the SEDs of galaxies with different galaxy types by means of spectral synthesis models. In this study we include Elliptical (E), Lenticular (S0), Spiral from Sa to Sd and Irregular (Irr) types.

Studies found that the observed properties of local field galaxies with the different galaxy types can be roughly matched by population with different SFRs. Therefore, using the BC03 software package we build 8 SFRs (corresponding to the types from E to Irr): a delta burst, a const star-formation system, and six μ -models with characteristic time-decays (exponentially decreasing SFR, $\tau = 1, 2, 3, 5, 10, 15$,

and 30 Gyr, see Eq. 2) chosen to match the sequence of colours from E-S0 to Sd (Bolzonella et al. 2000):

$$\psi(t) = [1 + \varepsilon M_{\text{PG}}(t)]\tau^{-1}\exp(-t/\tau) \quad (2)$$

where τ is the e-folding timescale, $M_{\text{PG}}(t) = [1 - \exp(-t/\tau)] - M_{\text{stars}} - M_{\text{remnants}}$ is the mass of gas that has been processed into stars and then returned to the interstellar medium (ISM) due to stellar evolution at time t , M_{stars} and M_{remnants} are the masses of stars and remnants at t , ε denotes the fraction of $M_{\text{PG}}(t)$ that can be recycled into new star formation.

By convolving these 8 SFRs with the SEDs of populations we generate 8 spectral families for a given spectral synthesis models. Each family (corresponding to a certain SFR) includes the SEDs at different ages.

In this study the number of ages (221, see Table 1) is reduced to 51 for each spectral family built with the GISSEL98 and BC03 models (refer to Bolzonella et al. 2000 for details), while it remains unchanged for the family built with the HPL07 and Yunnan models because the *Hyperz* code does not interpolate on the template grids and the template set must be densely populated (Maraston 2005).

2.3 Definition of Models

The observed spectra, provided by the *Hyperz* package, include 4 mean spectra of local E-, Sbc-, Scd- and Irr-type galaxies from Coleman et al. (1980, hereafter CWW, CWW-E, CWW-Sbc, CWW-Scd and CWW-Irr). These spectra extend from 1,400 to 10,000 Å originally and have been extended to the UV and near-IR regions by means of GISSEL98 spectra with parameters (SFR and age) selected to match the observed spectra at $z = 0$ (Bolzonella et al. 2000).

In all of the following computations, the template SED library is constituted by 8 theoretical spectral families (correspond to Burst, E, S0-Sd and Irr types) generated by spectral synthesis models and 4 observed galaxy spectra from CWW. For the sake of clarity 6 models are defined, each model differs from the other by changing the set of theoretical template SEDs (8 spectral families) in the *Hyperz* code: Models A/B/C/D use the theoretical template SEDs which were built with the GISSEL98/BC03/HPL07-s/Yunnan-s models (i.e., the models for SSPs), respectively, Models C2/D2 with the HPL07-b/Yunnan-b models (i.e., the models for BSPs). All the other relevant parameters introduced in the computations are fixed except the set of filters and the catalogue of galaxies. The description of parameters adopted by us is as follows.

- The combination of the CWW set of empirical SEDs with theoretical template spectra is used because the CWW set produces slightly better results than GISSEL98 (due to a Lyman blanketing effect) in the high redshift domain, and GISSEL98 models produce more accurate results than CWW templates alone in the low redshift domain. All 8 theoretical spectral families are used although there is no sensible gain when the set of 5 SFRs (1 delta burst, 3 μ -decaying, 1 constant star-formation system, Bolzonella et al. 2000) is used. Moreover, during the construction of theoretical template SEDs, we (i) assume that gas does not get recycled into new star formation once it has been processed into stars (i.e., $\varepsilon = 0$, see Eq. 2), (ii) neglect the attenu-

ation by dust, (iii) only use solar-metallicity EPS models because Bolzonella et al. (2000) found there is only a slight improvement on the accuracy of photo- z at $z \lesssim 1.5$ when several different metallicities are used together and found that including different metallicities does not affect the high redshift determinations, and, (iv) only use a certain IMF for each set of EPS models because that the photo- z estimates are approximatively the same, whatever the IMF used (Bolzonella et al. 2000).

- The reddening law of Calzetti et al. (2000) is used to dered the data because most of the fits to the galaxies from the Hubble Deep Field (HDF) by using the Calzetti's reddening law produce better χ^2 values than by using the other attenuation laws provided by the *Hyperz* package (Bolzonella et al. 2000). In addition, in the computations $A_V \equiv R_V \cdot E(B-V)$ ranges from 0. to 1.2 in steps of 0.2.

- The default values of flux decrements in the Lyman forest are used although their variation can affect the determination of photo- z , for example, decreasing them can induce an overestimate of photo- z .

- In the case of non-detection in filter i (characterized by the i -passband magnitude $m(i) = 99$ in the catalogue of galaxies), the contribution of the i -th filter is also taken into account in the χ^2 calculation by assuming the flux in the i -passband is 0 and the flux error equals to that deduced from the limiting magnitude ($\Delta F_{\text{obs}} = \Delta F_{\text{lim}}$). This assumption is the usual setting when one is dealing with a relatively deep survey in the considered filter (Bolzonella et al. 2000). The limiting magnitudes in the $F_{\text{UV}} N_{\text{UV}} ugriz$ passbands are set to 29, 29, 29, 30, 30, 30 and 30 mag, respectively.

- The minimum of photometric error Δm_{min} is set to 0.05 because there is no significant gain for $\Delta m_{\text{min}} \leq 0.05$ (about 5% accuracy, Bolzonella et al. 2000). When the input error from the catalogue is less than this value, Δm_{min} will replace the input error to avoid too small and non-realistic photometric errors.

- The set of cosmological parameters $(\Omega_M, \Omega_\Lambda, H_0) = (0.3, 0.7, 70)$ is used. The variation in this set of parameters [from (1.0, 0.0, 50) to (0.3, 0.7, 50)] only affects δ_z by less than 1% (Bolzonella et al. 2000), here H_0 in units of $\text{kms}^{-1} \text{Mpc}^{-1}$ and $\delta_z \equiv \sigma/(1 + \langle z \rangle)$, $\sigma = \sqrt{\sum_{i=1}^N |z_{\text{phot},i} - z_{\text{spec},i}|^2 / (N - 1)}$.

- The redshift ranges from 0 to 7 in a step of 0.02 instead of 0.05 (default value) although the primary z -step between 0.1 and 0.05 does not significantly affect the results (Bolzonella et al. 2000).

3 GALAXY SAMPLE

To check the ability of recovering photo- z s and morphological types of galaxies, we need to collect the galaxies with known spectroscopic redshifts and morphological classifications, and the redshift coverage of galaxies should be as wide as possible. But due to the limit of observations the galaxies with known classifications are relatively bright and their redshifts are low, two galaxy samples (termed spectroscopic and morphological) are used in this work, and they are used to study the ability of recovering photo- z s and galaxy types, respectively.

The galaxies in the spectroscopic sample are selected

Table 2. Characteristics of filters: the effective wavelength λ_{eff} and the surface of the normalized response function (width). Note the superscript in the first column denotes the value of airmass.

filters	$\lambda_{\text{eff}} [\text{\AA}]$	width $[\text{\AA}]$
F_{UV}	1539.781	252.031
N_{UV}	2313.893	729.648
u^0	3523.768	587.708
g^0	4798.005	1327.291
r^0	6258.645	1333.650
i^0	7655.315	1345.910
z^0	8984.980	1193.529




Figure 2. The spatial distribution of galaxies in the spectroscopic sample (red points, 6,531 galaxies). For comparison those mismatched galaxies between SDSS DR7 and GALEX DR4 in the *primate* spectroscopic sample, i.e., those SDSS DR7 galaxies not included in GALEX DR4, also are given (black points, 4,512 galaxies).




Figure 3. The transmission curves S_λ of F_{UV} , N_{UV} filters in GALEX system and $ugriz^0$ (for a null airmass) filters in SDSS system.

from the SDSS DR7. In order to ensure that the selected objects can be used in the studies of Section 4, we use the following criteria: the spectroscopic redshifts must be available (from *SDSS.SpecObj*), and GALAXY must be defined as the object class by all of the following three methods: spectral classification (*SDSS.SpecObj.specClass=2*), plPlugMap type (*SDSS.SpecObj.objType=0*) and morphological classification (*SDSS.Galaxy.type=3*). Initially we obtain a sample of 773,126 galaxies from the SDSS DR7. To save the computational time we use one-seventy of this galaxy sample as the *primitive* spectroscopic sample, which totally comprises 11,043 galaxies and in which the galaxies are randomly selected.

The galaxies in the morphological sample are from the catalogue of Fukugita et al. (2007), which contains 2,253 galaxies with *Petrosian* magnitude r_p brighter than 16 mag in the north equatorial stripe from the SDSS DR3. The morphological classification in the catalogue of Fukugita et al. (2007) is obtained by visual inspection of images in the g band. Removal the objects mismatched with the SDSS DR7 (the matching radius is 6 arcsec) produces a initial morphological sample of 1,792 galaxies. Note that the redshifts of galaxies in this catalogue are low (less than 0.2).

3.1 Matching the SDSS/DR7 sample with GALEX/DR4

We adopt the matching method of Agueros et al. (2005) and Obric et al. (2006) between the SDSS DR7 and the GALEX DR4, i.e., the matching radius between SDSS DR7 and GALEX DR4 is 6.0 arcsec, the nearest neighbor is taken as a true associate if its distance is smaller than the matching radius. When the nearest object has no simultaneous F_{UV} and N_{UV} detections, the nearer object (within the matching radius) with simultaneous F_{UV} and N_{UV} detections will be taken as its associate (361 galaxies for spectroscopic sample). If all objects within the matching radius have no simultaneous F_{UV} and N_{UV} detections, the missing magnitude of the nearest object will be replaced by the corresponding one of the nearer objects (within the matching radius).

At last, we obtain a final spectroscopic sample of 6,531 galaxies. In Fig. 2 we give the spatial distribution of these matched (i.e., the spectroscopic sample) and mismatched galaxies in the *primitive* spectroscopic sample, from it we see that the galaxies in the spectroscopic sample are indeed randomly distributed. Also we obtain a final morphological sample of 1,502 galaxies. The spectroscopic and morphological galaxy samples are used in the studies of Sections 4 and 5, respectively.

SDSS DR7 provides the values of various magnitudes (such as *devMag*, *expMag*, *modelMag*, *petroMag*, *psfMag*, etc). We choose SDSS *modelMag* (exactly, the shorthand alias for *modelMag*) of galaxies since the SDSS *C-model* magnitude (based on de Vaucouleurs and exponential fitting) is an adequate proxy to use as a universal magnitude for all types of objects and agrees with *Petrosian* magnitude excellently for galaxies. Moreover, the SDSS *C-model* magnitude is a reliable estimate of galaxy flux and can account for the effects of local seeing and thus is less dependent on local seeing variations (www.sdss.org/DR7/algorithm/photometry.html).




Figure 4. Comparisons between spectroscopic (z_{spec}) and photometric (z_{phot}) redshifts for the spectroscopic galaxy sample (see Section 3.2). Left panels are for only using optical ($ugriz$) photometry. Right panels are for using the combination of UV with optical photometry ($F_{\text{UV}} N_{\text{UV}} ugriz$). Top and bottom panels are for Models A and B, respectively. In each panel dot-dashed, dashed and solid lines correspond to $|\Delta z| = 0.2, 0.5$ and 1.0 . Dots and crosses denote galaxies with $z_{\text{phot}} \leq 1.0$ and $z_{\text{phot}} > 1.0$. Red, black, green and purple symbols denote galaxies with the photo- z probability $P(\chi^2) > 99, 99 \geq P(\chi^2) > 90, 90 \geq P(\chi^2) > 68, P(\chi^2) \leq 68$ (see Eq. 1), respectively.

3.2 Filters

The filters, involved in the computations, include those equipped in the SDSS system and those in the GALEX system. The transmission curves (S_λ) of these filters are presented in Fig. 3. The data about the SDSS filters for a null airmass ($ugriz^0$) are from the BC03 package. The characteristics of these filters (the effective wavelength λ_{eff} and the surface of the normalized response function [width]) are presented in Table 2.

4 UV PHOTOMETRY, BINARY INTERACTIONS AND POPULATION SYNTHESIS MODELS ON PHOTO-Z DETERMINATIONS

In this section we will use the *Hyperz* code to obtain photo- z estimates of galaxies in the spectroscopic sample, and discuss the effects of UV photometry, binary interactions and population synthesis models on photo- z determinations.

All parameters adopted by us in the *Hyperz* procedure (including the set of template spectra, reddening law and so on) have described in Section 2.3. In Appendix A we present the influence of these parameters on photo- z estimates of galaxies in the spectroscopic sample.




Figure 5. Similar to Fig. 4, but for Models C and C2 (top and bottom panels).




Figure 6. Similar to Fig. 4, but for Models D and D2 (top and bottom panels).

4.1 UV photometry on photo- z determinations

4.1.1 The effect of UV photometry

We obtain photo- z estimates (z_{phot}^V) of galaxies in the spectroscopic sample by only using SDSS $ugriz$ data and those ($z_{\text{phot}}^{\text{UVV}}$) by using the full set of $F_{\text{UV}} N_{\text{UV}} ugriz$ photometry for Models A-D2 (see Section 2.3).

In the left panels of Figs. 4-6, we give the comparisons between SDSS spectroscopic (z_{spec}) and photometric (z_{phot}^V) redshifts for Models A-D2, respectively. From them we see




Figure 7. Comparison between z_{phot}^V and $z_{\text{phot}}^{\text{UVV}}$ (based on Model A) for the spectroscopic galaxy sample. The dot-dashed, dashed and solid lines have the same meaning as in Fig. 4. The numbers in this figure correspond to the spectroscopic subsample (see Section 4.1.2). Arrows point to the location of galaxies in the 3rd subsample for the sake of clarity. Different symbols (squares, circles, stars and dots) and colours (red, black, green and purple) denote galaxies with different z_{phot}^V and $z_{\text{phot}}^{\text{UVV}}$ probability: $P(\chi^2) > 99$, $99 \geq P(\chi^2) > 90$, $90 \geq P(\chi^2) > 68$, $P(\chi^2) \leq 68$ (see Eq. 1), respectively.




Figure 8. The colour-colour diagrams for the spectroscopic galaxy sample. The galaxies in the 1-4 subsamples, which are divided based on Model A, are in black, red, green and blue, respectively. Squares, circles and stars denote galaxies with $F_{\text{UV}} ugriz$, $N_{\text{UV}} ugriz$ and $F_{\text{UV}} N_{\text{UV}} ugriz$ detections in the 3-4 subsamples, respectively.




Figure 9. Distribution of magnitudes (from left to right: F_{UV}^* , N_{UV}^* , u^* , g^* , r^* , i^* , z^*) along the wavelength for the 4 spectroscopic galaxy subsamples, which are divided based on Model A. Black dots, red dots, downarrows and uparrows represent the galaxies in the 1-4 subsamples, respectively. The horizontal lines represent the corresponding magnitude-errors, ΔMag . The length of line located in the upper left corner denotes $\Delta \text{Mag} = 1$. The galaxies with $F_{\text{UV}} N_{\text{UV}} ugriz$ detections in the 3rd and 4th subsamples are in green and purple, and those with $N_{\text{UV}} ugriz$ detections are in blue and grey. For the sake of clarity, the galaxies in the 2-4 subsamples are shifted rightwards.




Figure 10. Probability of photometric redshift $P(\chi^2)$ (see Eq. 1) as a function of redshift for an arbitrary galaxy in the 2nd (left panel), 3rd (middle panel) and 4th (right panel) subsamples based on Model A. In each panel black and red curves (the probability has been reduced by half for the sake of clarity) are for using $ugriz$ and $F_{\text{UV}} N_{\text{UV}} ugriz$ photometry. Red vertical line corresponds to the spectroscopic value. Also in each panel the magnitudes and their uncertainties, which are multiplied by 5, are plotted for the corresponding galaxy.

that for most of galaxies ($\sim 71.3\%$ for Model C) z_{phot}^V is less than 1.0 and the difference between spectroscopic and photometric redshifts $|\Delta z^V|$ ($\equiv |z_{\text{spec}} - z_{\text{phot}}^V|$) is less than 0.2, and for part of galaxies ($\sim 27.4\%$ for Model C) z_{phot}^V ranges from 1.8 to 2.8 but z_{spec} is less than 0.2 (these galaxies are located within the corresponding box of the left panels in Figs. 4-6).

In the right panels of Figs. 4-6 we give the comparisons between z_{spec} and $z_{\text{phot}}^{\text{UVV}}$ for Models A-D2. By comparing with the corresponding left-hand panel, we find that the number of galaxies with $z_{\text{phot}}^{\text{UVV}} < 1.0$ and $|\Delta z^{\text{UVV}}|$ ($\equiv$




Figure 11. the number of non-catastrophic identifications N (squares— $ugriz$; triangles— $F_{UV}N_{UV}ugriz$), mean deviation $\langle \Delta z \rangle$ (thin lines) and standard deviation σ_z (thick lines) [solid line— $ugriz$; dot-dashed line— $F_{UV}N_{UV}ugriz$] as a function of z_{spec} in removing catastrophic identifications for Models A-D2.




Figure 12. Similar to Fig. 11, but as a function of $g-r$ colour.

$|z_{\text{spec}} - z_{\text{phot}}^{\text{UVV}}| < 0.2$ is largely increased ($\sim 98.0\%$ for Model C), and that of galaxies with $1.8 \leq z_{\text{phot}}^{\text{UVV}} \leq 2.8$ and $z_{\text{spec}} \leq 0.2$ is significantly decreased ($\sim 0.18\%$ for Model C). That is to say, the inclusion of UV photometry would make the difference between photometric and spectroscopic redshifts ($|\Delta z|$) smaller (from 1.8-2.8 to less than 0.2) for these galaxies. When we only use optical data to obtain the photo- z estimates of these galaxies, we must be cautious. Then what are the characteristics of these galaxies?




Figure 13. Similar to Fig. 11, but as a function of r -magnitude.

4.1.2 which galaxies are affected?

According to the results about $z_{\text{phot}}^{\text{V}}$ and $z_{\text{phot}}^{\text{UVV}}$, we divide the spectroscopic galaxy sample into 4 subsamples for each model, the galaxies in the 1-4 subsamples satisfy the following conditions in turn:

- (i) both $z_{\text{phot}}^{\text{V}}$ and $z_{\text{phot}}^{\text{UVV}}$ are less than 1.0, and the differences, $|\Delta z^{\text{V}}|$ and $|\Delta z^{\text{UVV}}|$, are less than 0.2;
- (ii) $z_{\text{phot}}^{\text{V}} \geq 1.0$ and $z_{\text{phot}}^{\text{UVV}} \leq 1.0$;
- (iii) both $z_{\text{phot}}^{\text{V}}$ and $z_{\text{phot}}^{\text{UVV}}$ are greater than 1.0;
- (iv) $z_{\text{phot}}^{\text{V}} \leq 1.0$ and $z_{\text{phot}}^{\text{UVV}} \geq 1.0$.

This division means that the photo- z estimates of galaxies in the 1st subsample could be obtained reasonably in the case of only using optical $ugriz$ data, the photo- z errors of galaxies in the 2nd subsample could be decreased by the combination of UV with optical data, the photo- z estimates of galaxies in the 3rd subsample could not be obtained reasonably even if UV data is combined, and the photo- z estimates of galaxies in the 4th subsample would be erroneously identified if UV photometry is included. In Fig. 7 we give a comparison between $z_{\text{phot}}^{\text{V}}$ and $z_{\text{phot}}^{\text{UVV}}$ for Model A, the galaxies in the 1-4 subsamples occupy the lower left (except 80 galaxies, for which $0.2 < \Delta z < 0.5$), upper left, upper right and lower right regions, respectively.

To investigate the properties of galaxies in each subsample, according to the results of Model A, in Fig. 8 we present the distribution of galaxies in the diagrams of $(F_{UV}^* - N_{UV}^*)$.vs. $(g^* - r^*)$, $(F_{UV}^* - u^*)$.vs. $(g^* - r^*)$, $(N_{UV}^* - u^*)$.vs. $(g^* - r^*)$ and $(u^* - r^*)$.vs. $(g^* - r^*)$, in Fig. 9 present the magnitude $(F_{UV}^*, N_{UV}^*, u^*, g^*, r^*, i^* \text{ and } z^*)$ distribution of galaxies along the wavelength, and in Fig. 10 we give the photo- z probability $P(\chi^2)$ (see Eq. 1) as a function of redshift for an arbitrary galaxy in the 2nd, 3rd and 4th subsamples in both cases of using $ugriz$ and $F_{UV}N_{UV}ugriz$ photometry, and in each panel of Fig. 10 the magnitudes and magnitude-errors of the corresponding galaxy are also shown. Note that all of these magnitudes are corrected for foreground extinction by using the Schlegel et al. (1998) reddening maps, the extinction law of




Figure 14. Comparison between $z_{\text{phot}}^{\text{NUV}}$ and $z_{\text{phot}}^{\text{UV}}$ (top panel, based on Model B) and comparison between $z_{\text{phot}}^{\text{FUV}}$ and $z_{\text{phot}}^{\text{UV}}$ (bottom panel, based on Model B) for the spectroscopic galaxy sample. The lines, symbols and colours have the same meaning as in Fig. 7, but the y-axis represents $z_{\text{phot}}^{\text{NUV}}$ or $z_{\text{phot}}^{\text{FUV}}$, respectively.

Schlegel et al. (1998) for SDSS *ugriz* magnitudes and that of Cardelli, Clayton & Mathis (1989, hereafter CCM) for GALEX F_{UV} and N_{UV} magnitudes. By averaging the CCM's extinction law over GALEX F_{UV} and N_{UV} bandpasses, Wyder et al. (2005) obtained $A_{F_{\text{UV}}}/E(B-V) = 8.374$ and $A_{N_{\text{UV}}}/E(B-V) = 8.741$ (Donas et al. 2007). Meanwhile, all of these magnitudes are k -corrected by using SDSS spectroscopic redshift z_{spec} and the K -correction v4.1.4 code of Blanton & Roweis (2007), in which the distance moduli are obtained by assuming the cosmological parameters $(\Omega_M, \Omega_\Lambda) = (0.3, 0.7)$.

(i) From Fig. 8 we see that the galaxies in the 1st and 2nd subsamples are redder ($g^* - r^* \gtrsim 0.70$) and bluer ($g^* - r^* \lesssim 0.70$) ones, respectively. The main reason that the photo- z s of galaxies in the 2nd subsample are erroneously identified by using optical photometry is that bluer galaxies are often accompanied by star formation (Rich et al. 2005),




Figure 15. F_{UV} magnitude of a solar-metallicity population (normalized to $1 M_\odot$) for HPL07 (triangles) and Yunnan (squares) models. Open and solid symbols represent SSP and BSP, respectively. Also shown are the results of BC03 (solid black line+circles).




Figure 16. F_{UV} magnitude of a solar-metallicity galaxy ($1 M_\odot$) at $z = 0$. Solid circles, squares, up-triangles, left-triangles, right-triangles, down-triangles and stars symbols are for Burst, E, S0, Sa, Sb, Sc and Sd galaxies. The gray and coloured (black, red, green, blue, cyan, magenta and yellow correspond to Burst, E, S0, Sa-Sd types, in turn) symbols refer to using Yunnan-s and Yunnan-b models, respectively.

leading to a probability function $P(\chi^2)$ with significant secondary peak in the photo- z calculation (see the left panel of Fig. 10). The secondary peak (in the regime $1.8 \lesssim z \lesssim 2.8$) is explained by that the $4000 \text{ \AA}$ break, which is the most commonly used spectral feature for optical photo- z determination, is greatly reduced [making it more difficult to use as a redshift indicator (Niemack et al. 2009, and references therein)], and by that in the $1.8 \lesssim z \lesssim 2.8$ region the $4000 \text{ \AA}$ break goes beyond the wavelength coverage of *ugriz* filters [making it impossible to use as a redshift indicator




Figure 17. Evolutions of N (top panels), $|\langle\Delta z\rangle|$ (middle panels) and σ_z (bottom panels) as a function of z_{spec} for Models A-D2 in removing catastrophic identifications. Left and right panels are for using *ugriz* and $F_{\text{UV}}N_{\text{UV}}\textit{ugriz}$ photometry, respectively.




Figure 18. Similar to Fig. 17, but a function of $g-r$ colour.

]. When UV data is combined, the Lyman break, which is exhibited by all galaxies, would emerge in the UV ranges if $1.8 \lesssim z \lesssim 2.8$, this would largely decrease the probability of the solution in the high redshift domain.

(ii) the 3rd spectroscopic subsample (including 30 galaxies) is constituted of 9 galaxies with the full set of photometry and 21 galaxies with $N_{\text{UV}}\textit{ugriz}$ detections (one has wrong photometry). 8 of the 9 galaxies with $F_{\text{UV}}N_{\text{UV}}\textit{ugriz}$ detections and 15 of the 21 galaxies with $N_{\text{UV}}\textit{ugriz}$ detections have fainter u -light ($\gtrsim 20.5$ mag) and larger u -magnitude errors (see Fig. 9 and the middle panel of Fig. 10), and therefore have bluer $F_{\text{UV}} - u^*$ and $N_{\text{UV}}^* - u^*$ colours and redder $u^* - \text{optical}^*$ colours (such as $u^* - g^*$ colour, see Fig. 8). The larger u -magnitude error leads to a smaller contribution of the u -band photometry to the χ^2 calculation (see Eq. 1), the probability function has two peaks for degeneracy of the fit parameters if only optical




Figure 19. Similar to Fig. 17, but a function of r -magnitude.




Figure 20. The percent difference in the number of non-catastrophic identifications $dN \equiv (N_{x2} - N_x)/N_{x2}$, $x=C$ or D) between Models C and C2 (squares) and between Models D and D2 (triangles) as a function of z_{spec} (Left panels), $g-r$ colour (middle panels) and r -magnitude (right panels). Upper and bottom panels are for using *ugriz* and $F_{\text{UV}}N_{\text{UV}}\textit{ugriz}$ photometry, respectively.

photometry is used. Even if the UV photometry is included, the high redshift resolution (at $z > 2.3$, see Fig. 7 and the middle panel of Fig. 10) could not be excluded by the *Hyperz* code. This is explained by that the Lyman break would move to wavelength longer than $3000 \text{ \AA}$ [i.e., emerge in the u band] and the $4000 \text{ \AA}$ break would go beyond the wavelength coverage of *ugriz* filters. Note that the confidence levels are very low in the determinations of both z_{phot}^V and $z_{\text{phot}}^{\text{UVV}}$ for most of galaxies in this subsample (refer to Fig. 7).

(iii) the 4th subsample (9 galaxies) is constituted of 2 galaxies with the full set of photometry and 7 galaxies with $N_{\text{UV}}\textit{ugriz}$ detections. The galaxies in this subsample have fainter N_{UV} light (see Fig. 9 and the right panel of Fig. 10), therefore have redder $N_{\text{UV}}^* - \text{optical}^*$ colours (such as $N_{\text{UV}}^* - u^*$ colour, see Fig. 8). Because the fainter N_{UV} light also can

be produced by high redshift galaxies, for which the Lyman break would move to the N_{UV} passband, the *Hyperz* code is unable to decide between the high redshift and low redshift solutions. From the right panel of Fig. 10 we see that the $P(\chi^2)$ of z_{phot}^V is higher than that of z_{phot}^{UVV} .

In summary, the addition of UV photometry would increase the number of non-catastrophic (i.e., $|\Delta z| = |z_{\text{spec}} - z_{\text{phot}}| < 1.0$) identifications, and decrease the mean deviation $\langle \Delta z \rangle$ ($\equiv \Sigma \Delta z / N$) and standard deviation σ_z ($\equiv \sqrt{\Sigma (\Delta z - \langle \Delta z \rangle)^2 / (N - 1)}$) when all galaxies are included. And, the 2nd spectroscopic subsample of galaxies, for which the photo- z errors are largely decreased by the inclusion of UV data, mainly locates in the redshift $z_{\text{spec}} < 0.2$ and the colour $g^* - r^* \lesssim 0.7$ regions, and has not significant difference in the UV and optical magnitudes in comparison with the 1st subsample. Therefore, if the galaxy is a bluer one, we had better combine UV with optical photometry to obtain its photo- z estimate. If the N_{UV} light of a galaxy is too faint, its photo- z estimate would be identified erroneously when UV data is included, and if the u -light of a galaxy is too faint and the u -magnitude error is too large, its photo- z estimate still would be erroneously identified even if UV photometry is included in the *Hyperz* code.

4.1.3 Dependence of photo- z accuracy

To investigate the performance of photo- z estimate in removing catastrophic identifications, in Tables B1-B3 we give the number of non-catastrophic identifications N , mean deviation $\langle \Delta z \rangle$ and standard deviation σ_z as a function of spectroscopic redshift z_{spec} , $g-r$ colour and r -magnitude for Models A-D2. Each table includes the cases of using *ugriz* and $F_{UV}N_{UV}$ *ugriz* photometry. The results in Tables B1-B3 are plotted in Figs. 11-13, respectively.

Fig. 11 gives the evolutions of N , $\langle \Delta z \rangle$ and σ_z with z_{spec} for Models A-D2.

- First, from the evolution of the mean deviation $\langle \Delta z \rangle$ with z_{spec} we see that if *ugriz* photometry is used the mean photometric redshift $\langle z_{\text{phot}} \rangle$ ($\equiv \Sigma z_{\text{phot}} / N$) in the regions of $z_{\text{spec}} \lesssim 0.05$ and $0.25 \lesssim z_{\text{spec}} \lesssim 0.45$ is greater (~ 0.05 at the most), and that in the region of $0.05 \lesssim z_{\text{spec}} \lesssim 0.25$ (~ 0.02 at the most) is less than the mean spectroscopic redshift $\langle z_{\text{spec}} \rangle$ ($\equiv \Sigma z_{\text{spec}} / N$) for Models A, B, D and D2. For Models C and C2, $\langle z_{\text{phot}} \rangle$ is greater than $\langle z_{\text{spec}} \rangle$ at all redshift bins. The maximal $|\langle \Delta z \rangle|$ locates in the regions of $z_{\text{spec}} \lesssim 0.05$ and $0.25 \lesssim z_{\text{spec}} \lesssim 0.45$ for all models. The addition of UV data almost does not vary the mean deviation except in the region $0.5 \lesssim z_{\text{spec}} \lesssim 0.6$, in which the number of galaxies is smaller.

- Secondly, from the correlation between σ_z and z_{spec} we see that the standard deviation σ_z in the $z_{\text{spec}} \lesssim 0.3$ region is less than that in the $z_{\text{spec}} \gtrsim 0.3$ region if only *ugriz* data is used, the addition of UV data decreases σ_z in the region of $z_{\text{spec}} \gtrsim 0.3$ for all models.

- At last, from Fig. 11 we see that the addition of UV photometry significantly increases the number of non-catastrophic identifications in the low redshift domain for all models.

Fig. 12 presents the evolutions of N , $\langle \Delta z \rangle$ and σ_z as a

function of $g-r$ colour, reveals the dependence of photo- z accuracy on $g-r$ colour.

- First, from the evolution of $\langle \Delta z \rangle$ with $g-r$ colour, we see that in the case of only using *ugriz* photometry the mean deviation $\langle \Delta z \rangle$ is positive in the $0.8 \lesssim g-r \lesssim 1.4$ region, and in the $0.3 \lesssim g-r \lesssim 0.8$ region $\langle \Delta z \rangle$ is negative and $|\langle \Delta z \rangle|$ increases with decreasing $g-r$ colour for Models A, B, D and D2. For Models C and C2 the mean deviation $\langle \Delta z \rangle$ is less than zero at all $g-r$ color bins and $\langle \Delta z \rangle$ almost does not vary with $g-r$ colour. The inclusion of UV data can decrease $|\langle \Delta z \rangle|$ in the $0.3 \lesssim g-r \lesssim 0.8$ region for Models A, B, D and D2, while increase $|\langle \Delta z \rangle|$ at the redder $g-r$ colour for Models C and C2.

- Secondly, from the evolution of σ_z with $g-r$ colour, we see that the standard deviation in the $0.8 \lesssim g-r \lesssim 1.2$ region is less than that in the other regions if *ugriz* photometry is used, and the inclusion of UV data would decrease the standard deviation in the regime $0.3 \lesssim g-r \lesssim 0.6$ (except for Model B) for all models.

- At last, from the correlation between N and $g-r$ colour, we see that the addition of UV photometry increases the number of non-catastrophic identifications in the region $g-r \lesssim 0.8$.

From Fig. 13, which gives the evolutions of $\langle \Delta z \rangle$, σ_z and N as a function of r -magnitude, we see that the mean deviation $\langle \Delta z \rangle$ and standard deviation σ_z are independent of r -magnitude in both cases of using *ugriz* and $F_{UV}N_{UV}$ *ugriz* photometry, the inclusion of UV photometry increases the number of non-catastrophic identifications in the range $17.0 \lesssim r \lesssim 18.5$, and increases σ_z for all models.

In summary, if catastrophic identifications are removed, the inclusion of UV data raises the number of non-catastrophic identifications in the low redshift, $g-r \lesssim 0.8$ and $17 \leq r \leq 18.5$ regions, decreases the mean deviation in the high redshift ($z_{\text{spec}} \gtrsim 0.5$) and $0.3 \lesssim g-r \lesssim 0.8$ regions, and also decreases the standard deviation in the $z_{\text{spec}} \gtrsim 0.3$ and $0.3 \lesssim g-r \lesssim 0.6$ regions.

4.1.4 F_{UV} or N_{UV} photometry on photo- z determinations

If we only combine F_{UV} or N_{UV} with optical photometry, what results would happen? Using the set of N_{UV} *ugriz* photometry, we obtain photo- z estimates ($z_{\text{phot}}^{N_{UV}V}$) of galaxies in the spectroscopic sample for Models A-D2, then compare them with the results obtained by using the set of $F_{UV}N_{UV}$ *ugriz* photometry (i.e., z_{phot}^{UVV}). For the sake of size, in the top panel of Fig. 14 we only give a comparison between z_{phot}^{UVV} and $z_{\text{phot}}^{N_{UV}V}$ for Model B. From it we see that for most of galaxies $z_{\text{phot}}^{N_{UV}V}$ is lower than 1.0 and $z_{\text{phot}}^{N_{UV}V} > z_{\text{phot}}^{UVV}$. That is to say, only N_{UV} is combined with *ugriz* photometry, the derived photo- z s are reasonable for most of galaxies, the number of non-catastrophic identifications N is increased, but the mean deviation and standard deviations are raised in comparison with those in the case of using $F_{UV}N_{UV}$ *ugriz* photometry (due to $z_{\text{phot}}^{N_{UV}V} > z_{\text{phot}}^{UVV}$). The reason that the inclusion of N_{UV} light decreases the probability of the solution in the region of $1.8 \lesssim z \lesssim 2.4$ is that the Lyman break would emerge in the N_{UV} passband. The overestimation of photo- z is explained by that the main

spectral feature still is the 4000 Å break, which is unable to break the degeneracy among the fit parameters if $z \lesssim 0.6$.

Using the set of $F_{UV}ugriz$ photometry, we also obtain photo- z estimates ($z_{phot}^{F_{UV}}$) for the spectroscopic galaxy sample. In the bottom panel of Fig. 14 we give a comparison between z_{phot}^{UVV} and $z_{phot}^{F_{UV}}$ for Model B. From it we see that $z_{phot}^{F_{UV}}$ is close to z_{spec} . The agreement is caused by the fact that the Lyman break would emerge in the F_{UV} passband if the redshift is low. Above conclusions are also true for the other Models.

In Tables B1-B3 we also give N , $\langle\Delta z\rangle$ and σ_z of Model B as a function of redshift, $g-r$ and r -magnitude when using $N_{UV}ugriz$ or $F_{UV}ugriz$ photometry. From them we see that the catastrophic objects mainly are those galaxies with $z_{spec} \gtrsim 0.3$, $g-r \gtrsim 0.8$ and $r \gtrsim 18$ mag when N_{UV} photometry is combined with optical photometry. Also, from them we see the catastrophic objects are those galaxies with $z_{spec} \lesssim 0.15$, $0.4 \lesssim g-r \lesssim 1.0$ and $16.5 \lesssim r \lesssim 18$ mag if F_{UV} photometry is combined with optical data. That is to say, only including N_{UV} or F_{UV} photometry would cause to erroneous photo- z estimates for those faint 'redder' high-redshift galaxies and those 'bluer' low-redshift (intermediate r -magnitude) galaxies, respectively. The number of catastrophic identifications in the case of using $F_{UV}ugriz$ photometry is less than that of using $N_{UV}ugriz$ photometry.

4.2 Binary interactions on photo- z determinations

Binary interactions would significantly change the UV spectra of populations at age $t \gtrsim 1$ Gyr (see Fig. 1). Using the HPL07-b, HPL07-s, Yunnan-b and Yunnan-s population synthesis models we get the F_{UV} , N_{UV} , u , g , r , i and z magnitudes of a solar-metallicity population with a mass of $1 M_{\odot}$. By comparison we find the differences in the N_{UV} , u , g , r , i and z magnitudes among these models are slight, while the difference in the F_{UV} magnitude caused by binary interactions, i.e., that between HPL07-s and HPL07-b or that between Yunnan-s and Yunnan-b models, is significant at $\log(t/yr) \gtrsim 9.0$ (see Fig. 15). Then whether binary interactions would affect photo- z estimates?

Comparing the photo- z estimates of galaxies in the spectroscopic sample between Models C and C2 (Fig. 5), and those between Models D and D2 (Fig. 6), we find that binary interactions do not systematically affect photo- z determinations. One reason for this is that binary interactions vary significantly the SEDs of populations (or Burst-type galaxies) only at late ages ($t \gtrsim 1$ Gyr), leading that at redshift $z = 0$ the difference in the SEDs, F_{UV} , N_{UV} , u , g , r , i and z magnitudes of non-Burst galaxies between Models C and C2 and that between Models D and D2 is slight at a given age and galaxy type. In Fig. 16 we give the F_{UV} magnitude of Burst, E, S0 and Sa-Sd galaxies at redshift $z = 0$ by using the Yunnan-s and Yunnan-b models. Even if at redshift $z = 0.6$, binary interactions only make the ISEDs of populations (or Burst-type galaxies) bluer in the regime $\lambda < 4000$ Å according to $z = (\lambda' - \lambda_0)/\lambda_0$ and $\lambda_0 = 2500$ Å, beyond which the effect of binary interactions on the ISEDs is slight. These bluer UV and near-optical ISEDs can be easily offset by the decrements of visual extinction A_V and age, absence of the Lyman break and the adoption of the different reddening laws during the fitting process.

In Figs. 17-19 (also generated from Tables B1-B3 in turn), we present N , the absolute value of mean deviation $|\langle\Delta z\rangle|$ and σ_z as a function of z_{spec} , $g-r$ colour and r -magnitude for Models A-D2 in both cases of using $ugriz$ and $F_{UV}N_{UV}ugriz$ photometry, also the catastrophic identifications are removed. From Figs. 17-19 we see that binary interactions decrease and increase $|\langle\Delta z\rangle|$ in the regions of $0.3 \lesssim g-r \lesssim 0.8$ and $r \gtrsim 17.8$ mag in the case of using optical photometry, respectively (see the comparisons between Models D and D2 in Figs. 18 and 19), and decrease the standard deviation σ_z at all r magnitudes for both cases of using $ugriz$ and $F_{UV}N_{UV}ugriz$ data (see Fig. 19).

In Fig. 20 we present the percent difference in the number of non-catastrophic identifications between Models C and C2 and that between Models D and D2 ($dN \equiv (N_{x2} - N_x)/N_{x2}$, $x=C$ or D) as a function of z_{spec} , $g-r$ colour and r -magnitude. From it we see that N_{D2} is less than N_D at $0.2 \lesssim z_{spec} \lesssim 0.3$ in both cases of using $ugriz$ ($\sim 10\%$ at the most) and $F_{UV}N_{UV}ugriz$ ($\sim 2\%$ at the most) photometry, greater than N_D in the region $0.3 \lesssim g-r \lesssim 0.8$ when only using optical data ($\sim 23\%$ at the most), and almost equal to N_D in the whole $g-r$ region (except $g-r \gtrsim 1.2$ mag, in which N is smaller) when using the full set of photometry.

In brief, the inclusion of binary interactions mainly affects the photo- z determinations in the case of only using optical photometry, in an exact word, it increases the number of non-catastrophic identifications and decreases the deviations in the region of $0.3 \leq g-r \leq 0.8$ mag if only optical photometry is used.

4.3 EPS models on photo- z determinations

First, from the evolutions of the number of non-catastrophic identifications N with z_{phot} , $g-r$ colour and r -magnitude in the top panels of Figs. 17-19, we see that the difference in N among models A-D2 is smaller if $F_{UV}N_{UV}ugriz$ data is used, and see that N decreases from Model B, Model A, Models D/D2 to C/C2 (i.e., $N_B > N_A > N_{D/D2} > N_{C/C2}$) in the regions of $0.05 \lesssim z_{spec} \lesssim 0.15$, $0.3 \lesssim g-r \lesssim 0.8$ and $17 \lesssim r \lesssim 18.2$ mag when only optical data is used.

Secondly, from the middle and bottom panels of Figs. 17, we see that (i) the absolute value of mean deviation $|\langle\Delta z\rangle|$ and standard deviation σ_z of Models C/C2 are greater than those of the other models in the low redshift domain for both cases of using $ugriz$ and $F_{UV}N_{UV}ugriz$ photometry; (ii) among Models A, B, D and D2, the difference of $|\langle\Delta z\rangle|$ in the regions of $z_{spec} \lesssim 0.05$ and $z_{spec} \gtrsim 0.5$ and the difference of σ_z in the region of $z_{spec} \gtrsim 0.5$ are larger (except σ_z in the case of using $F_{UV}N_{UV}ugriz$ data), furthermore, in the $0.1 \lesssim z_{spec} \lesssim 0.5$ region the $|\langle\Delta z\rangle|$ and σ_z of Model B are greater than those of Models A, D and D2 for both cases of using $ugriz$ and $F_{UV}N_{UV}ugriz$ photometry (except σ_z by using optical data in the $0.3 \lesssim z_{spec} \lesssim 0.4$ region); (iii) the evolutions of $|\langle\Delta z\rangle|$ and σ_z with redshift of Models D and D2 are similar to those of Model A.

Moreover, from the middle and bottom panels of Figs. 18, we see that in the $g-r \lesssim 0.7$ region the $|\langle\Delta z\rangle|$ and σ_z of these models (except σ_z in the case of using $F_{UV}N_{UV}ugriz$ data) are significantly different, and that the $|\langle\Delta z\rangle|$ and σ_z of Model B are significantly lower than those of the other models for both cases of using $ugriz$ and $F_{UV}N_{UV}ugriz$ photometry.

Table 3. Morphological index T of Fukugita et al. (2007) and Ours.

galaxy Type	$T(\text{Fuk})$	$T(\text{Ours})$
Unclassified	-1	...
Burst	...	1
E	0	2
S0	1	3
Sa	2	4
Sb	3	5
Sc	4	6
Sd	5	7
Irr	6	8
CWW-E	...	9
CWW-Sbc	...	10
CWW-Scd	...	11
CWW-Irr	...	12

At last, from the middle and bottom panels of Figs. 19, we see that the mean and standard deviations of Models C and C2 are greater than those of the other models in both cases of using *ugriz* and $F_{\text{UV}}N_{\text{UV}}\textit{ugriz}$ photometry, and in the intermediate r -magnitude region the $|\langle\Delta z\rangle|$ of Models A, D and D2 are lower in both cases of using *ugriz* and $F_{\text{UV}}N_{\text{UV}}\textit{ugriz}$ photometry.

From above comparisons, we see that in the case of using $F_{\text{UV}}N_{\text{UV}}\textit{ugriz}$ photometry the number of non-catastrophic identifications is similar for all models, and in the case of using *ugriz* photometry the number of non-catastrophic identifications of Models B is greater than that of the other models in the low redshift, bluer $g-r$ and faint r -magnitude regions. In both cases of using *ugriz* and $F_{\text{UV}}N_{\text{UV}}\textit{ugriz}$ photometry, the mean and standard deviations of Model B are lower at $g-r \lesssim 0.7$, and $|\langle\Delta z\rangle|$ of Models A, D and D2 are relatively lower at almost all redshift bins and at intermediate r magnitudes.

5 UV PHOTOMETRY AND BINARY INTERACTIONS ON GALAXY MORPHOLOGY

Bolzonella et al. (2000) claimed that the *Hyperz* code allows to obtain photo- z estimate and the best fit parameters across the whole space of galaxy. Because of the lack of spectral resolution, only the rough spectral types could be retrieved from broad-band photometry, in an exact word, early types could be reliably and easily identified while late types are usually misidentified because of the degeneracy between spectral type, age of the stellar population and visual extinction.

Using the *Hyperz* code and the morphological sample of 1,502 galaxies, which are from the catalogue of Fukugita et al. (2007) and have matched with the SDSS DR7 and GALEX DR4 (see the text in Section 3 for details), we recover their morphological types by using *ugriz* and $F_{\text{UV}}N_{\text{UV}}\textit{ugriz}$ photometry for Models A-D2, respectively. The corresponding galaxy types of morphological index of Fukugita et al. (2007), $T(\text{Fuk})$, and ours, $T(\text{Ours})$, are presented in Table 3. Note that half-integer classes are allowed for $T(\text{Fuk})$.

In Figs. 21-22 we give the histograms of r -magnitude




Figure 21. Histogram of r -magnitude distribution for E (thick black solid line), S0 (thick red dashed), Sa (thick green dot-dashed), Sb (thick blue dotted), Sc (thick magenta dash-dot-dot), Sd (thin grey solid) and Irr (thin grey dashed) morphological subsample (see the text in Section 3). Note that the half-integer classes are designated as the corresponding integer classes in this figure.




Figure 22. Similar to Fig. 21, but for spectroscopic redshift z_{spec} . Left panel is for E, S0 and Sa types, right panel for Sb, Sc, Sd and Irr types. The lines have the same meaning as in Fig. 21.




Figure 23. Histogram of morphology-index $T(\text{Fuk})$ distribution for morphological galaxy sample. Grey solid line is for the original catalogue of Fukugita et al. (2007), red dotted line is for the sample of galaxies matched with the SDSS DR7 and GALEX DR4, also shown are the sample of galaxies only matched with the SDSS DR7 (black solid line). The corresponding galaxy types of $T(\text{Fuk})$ are given in Table 3.

Table 4. Reliability and completeness for early- and late-type selection by the *Hyperz* code (including Models A-D2). The upper and lower parts correspond to using *ugriz* and $F_{UV}N_{UV}ugriz$ photometry, respectively. This table is based on the morphology sample of 1,502 galaxies.

Model	early type		late type	
	reliability	completeness	reliability	completeness
<i>ugriz</i>				
A	0.74	0.60	0.62	0.54
B	0.69	0.60	0.58	0.46
C	0.83	0.39	0.54	0.73
C2	0.85	0.40	0.54	0.74
D	0.72	0.58	0.60	0.53
D2	0.74	0.63	0.65	0.52
$F_{UV}N_{UV}ugriz$				
A	0.70	0.58	0.57	0.49
B	0.70	0.58	0.57	0.49
C	0.86	0.43	0.57	0.74
C2	0.87	0.57	0.66	0.72
D	0.71	0.66	0.65	0.47
D2	0.71	0.67	0.66	0.45

Table 5. Reliability and completeness for selecting early- and late-type galaxies by the three morphology selection parameters: the concentration index C , profile likelihood and $u - r$ colour.

Selection Rule	reliability	completeness
early type		
$C > 2.6$	0.83	0.74
$P_{deV} > P_{exp}$	0.76	0.77
$u - r \geq 2.22$	0.71	0.75
late type		
$C < 2.6$	0.82	0.62
$P_{deV} < P_{exp}$	0.85	0.49
$u - r \leq 2.22$	0.77	0.40

and spectroscopic redshift z_{spec} distributions for E, S0, Sa, Sd and Irr subsamples, respectively, and in Fig. 23 give the histogram of $T(Fuk)$ distribution for the morphological sample. From them we see that the galaxies in the morphological sample are relatively bright ($r < 16$.mag, see Fig. 21) and their redshifts are less than 0.2 (see Fig. 22), and that most

Table 6. Similar to Table. 5, but for the four new constructed colour selection criteria.

Selection Rule	reliability	completeness
early type		
$N_{UV} - u \geq 1.94$	0.89	0.65
$F_{UV} - u \geq 3.16$	0.88	0.37
$5.27 - 1.30(u - r) > (N_{UV} - u)$	0.88	0.65
$5.77 - 1.47(u - r) > (F_{UV} - u)$	0.89	0.64
late type		
$N_{UV} - u < 1.94$	0.73	0.73
$F_{UV} - u < 3.16$	0.54	0.77
$5.27 - 1.30(u - r) < (N_{UV} - u)$	0.73	0.72
$5.77 - 1.47(u - r) < (F_{UV} - u)$	0.73	0.73




Figure 24. Comparisons in the morphological index of galaxies between ours and those of Fukugita et al. for Models A-D2. Black and red squares represent using *ugriz* and $F_{UV}N_{UV}ugriz$ photometry, respectively, and their area represents the number of galaxies on the grid. The area of square located in the upper left corner of each panel denotes the number of galaxies is 50. Crosses show the correlation between $T(Ours)$ and $T(Fuk)$.

of them belong to early types, Sd and Irr types are scarce (see Fig. 23).

5.1 UV photometry on morphological types

In Fig. 24 we compare the retrieved morphological index $T(Ours)$ with those of Fukugita et al. (2007), $T(Fuk)$, for Models A-D2, note that those galaxies with $|z_{phot} - z_{spec}| > 1$ are excluded.

From the comparisons in Fig. 24 we indeed see that in the case of only using optical photometry early types ($T(Fuk) < 2.0$) can be easily retrieved and late types ($T(Fuk) \geq 2.0$) are often misidentified as early types for all models except for Models C-C2. For Models C-C2 both early and late types can be reliably identified. And, if $F_{UV}N_{UV}ugriz$ photometry is used the situation has not changed significantly.

Moreover, from Fig. 25, which gives the difference in the number of galaxies between in the cases of using *ugriz* and $F_{UV}N_{UV}ugriz$ photometry on the plane of $T(Ours)$.vs. $T(Fuk)$, we see that the probability that early types ($T(Fuk) \leq 2.0$) are classified as Burst and E (and CWW-E for Models A-B) types is decreased and in-




Figure 25. The difference in the number of galaxies between in the case of using $ugriz$ [$N(ugriz)$] and in the case of using $F_{UV}N_{UV}ugriz$ photometry [$N(F_{UV}N_{UV}ugriz)$] on the plane of $T(\text{Ours})$.vs. $T(\text{Fuk})$ for Models A-D2. Black and red symbols represent the difference $dN \equiv N(ugriz) - N(F_{UV}N_{UV}ugriz)$ is positive and negative, respectively. Also, The size of squares scales with the number of galaxies, and the area of square located in the upper left corner of each panel also denotes the number is 50.




Figure 26. Similar to Fig. 25, but for the difference in the number of galaxies between Models D and D2 on the plane of $T(\text{Ours})$.vs. $T(\text{Fuk})$ (i.e., between neglecting and including binary interactions). Left and right panels are for using $ugriz$ and $F_{UV}N_{UV}ugriz$ photometry, respectively. Red and black squares denote that the number of galaxies of Model D on the grid is greater and less than that of Model D2, respectively. The size of symbols has the same meaning as in Fig. 25.




Figure 27. Distribution of galaxies in colour-colour diagrams (black, red, green, blue, cyan, yellow and magenta symbols denote E, S0, Sa-Sd and Irr types, respectively). Also shown are the lines corresponding to the four criteria: $F_{UV} - u = 3.16$ (horizontal line, top panel), $F_{UV} - u = 5.77 - 1.47(u - r)$ (oblique line, top panel), $N_{UV} - u = 1.94$ (horizontal line, bottom panel) and $N_{UV} - u = 5.27 - 1.30(u - r)$ (oblique line, bottom panel).

creased, respectively, and that the probability that late types ($T(\text{Fuk}) > 2.0$) are classified as CWW-Sbc and CWW-Scd types is increased.

To investigate the effect of UV photometry on the determinations of morphological types and the performance of the retrieved morphology by the *Hyperz* code, we define $\langle \Delta T_i \rangle \equiv \sum_{k=1}^{N_i} [T(\text{Ours}) - T(\text{Fuk})] / N_i$ and $\sigma_{T_i} \equiv \sqrt{\sum_{k=1}^{N_i} [T(\text{Ours}) - T(\text{Fuk}) - \langle \Delta T_i \rangle]^2 / (N_i - 1)}$, in which N_i is the number of non-catastrophic objects (i.e., $|z_{\text{spec}} - z_{\text{phot}}| < 1$.) classified as i -type by Fukugita et al. (2007, i.e., $T(\text{Fuk}) = i$) and i is allowed to be half-integer. In Figs. C1-C3 we present the evolutions of $|\langle \Delta T_i \rangle|$, σ_{T_i} and N_i as a function of spectroscopic redshift z_{spec} , $g - r$ colour and r -magnitude for Models A-D2.

The following comparisons in $|\langle \Delta T_i \rangle|$ and σ_{T_i} are based on those bins with $N_i \geq 10$ because the small value in N_i would lead to a larger error in $|\langle \Delta T_i \rangle|$ and σ_{T_i} . In each panel of Figs. C1-C3 the two vertical lines indicate the place of $N_i = 10$, therefore we compare $|\langle \Delta T_i \rangle|$ and σ_{T_i} in the region enclosed by two horizontal lines (shaded region of Figs. C1-C3).

- First, from Fig. C1, we see that the inclusion of UV light increases N_i of all models for $T(\text{Fuk}) \geq 1.0$ [the increment increases with $T(\text{Fuk})$], decreases $|\langle \Delta T_i \rangle|$ and σ_{T_i} of Models C and C2 for $T(\text{Fuk}) = 0.0 - 0.5$ at all redshift bins

and for $1.0 \leq T(\text{Fuk}) \leq 3.0$ (except = 2.5) at $z_{\text{spec}} \lesssim 0.04$, and also decreases $|\langle \Delta T_i \rangle|$ and σ_{T_i} of Models A and B for $T(\text{Fuk})=2.0-2.5$ at all redshift bins. Moreover, we also see that all models have the similar trends of $|\langle \Delta T_i \rangle|$ and σ_{T_i} with redshift for a given $T(\text{Fuk})$ when the full set of photometry is used: both $|\langle \Delta T_i \rangle|$ and σ_{T_i} are independent of z_{spec} for all galaxy types, and $|\langle \Delta T_i \rangle|$ increases with $T(\text{Fuk})$ [except Models C and C2, for which $|\langle \Delta T_i \rangle|$ is independent of $T(\text{Fuk})$].

- Secondly, from Fig. C2, we see that the inclusion of UV data increases N_i of all models for $2.5 \leq T(\text{Fuk}) \leq 4.0$ in the blue $g-r$ region [correspondingly, the left vertical line moves bluewards significantly, and N_i increases with $T(\text{Fuk})$], decreases $|\langle \Delta T_i \rangle|$ and σ_{T_i} of Models C and C2 at $T(\text{Fuk}) \leq 2.5$ and decreases $|\langle \Delta T_i \rangle|$ and σ_{T_i} of all models at $T(\text{Fuk})=3.0$ in the region of $g-r \lesssim 0.8$ mag. Moreover, if UV data is used, $|\langle \Delta T_i \rangle|$ correlates with $g-r$ colour and $T(\text{Fuk})$: $|\langle \Delta T_i \rangle|$ increases with redder $g-r$ ($\gtrsim 0.8$) colour for $T(\text{Fuk}) \gtrsim 2.0$ (i.e., late types) and increases with $T(\text{Fuk})$ for all models except Models C and C2.

- At last, from Fig. C3, we see that the inclusion of UV data increases N_i of all models for late types in the region of $r \gtrsim 14$ mag, decreases $|\langle \Delta T_i \rangle|$ and σ_{T_i} of Models C and C2 for $T(\text{Fuk}) \leq 1.0$ at all r magnitudes, and decreases σ_{T_i} and increases $|\langle \Delta T_i \rangle|$ of Models A and B at $2.0 \leq T(\text{Fuk}) \leq 3.5$ at almost all r magnitudes. The standard deviation σ_{T_i} is relatively large for fainter ($r \gtrsim 16$ mag) galaxies in the case of using *ugriz* data, when UV photometry is included σ_{T_i} almost does not vary with increasing r -magnitude except in the region of $r \gtrsim 16$ mag.

5.2 Binary interactions on morphological types

To obtain the influence of binary interactions on the retrieved morphological index, in Fig. 26 we give a difference in the number of non-catastrophic objects between Models D and D2 on the plane of $T(\text{Ours})$ and $T(\text{Fuk})$ for *ugriz* and $F_{\text{UV}}N_{\text{UV}}$ *ugriz* photometry.

From Figs. 24 and 26, we see that the inclusion of binary interactions mainly affects the determinations of E and S0 types in both cases of using optical and the full set of photometry, and the difference in the case of using optical photometry is more significant than that in the case of using $F_{\text{UV}}N_{\text{UV}}$ *ugriz* photometry. This is explained by that binary interactions make the UV flux of Burst-type galaxies higher only at $\log(t/\text{yr}) \gtrsim 9$, and these bluer UV spectra can be offset by the variation of other parameters (such as the decrements of age and visual extinction, etc) during the fitting process (also see the discussion in Section 4.2).

5.3 Comparison with other morphology selection criteria

In this section we will give a comparison in the performance of selecting early- (E, S0 and Sa) and late-type (Sb, Sc, Sd and Irr) galaxies between by using the *Hyperz* code (this work) and by using the three morphology-sensitive parameters (the concentration index C , the profile likelihood and $u-r$ colour). The concentration index C is defined as the ratio of the radii containing 90% and 50% of the *Petrosian* r galaxy light, i.e., $C \equiv r_{90}/r_{50}$. The profile likelihood is

defined as the difference between the deVaucouleur's (P_{dev}) and exponential (P_{exp}) profile likelihoods in the r band.

In Tables. 4 and 5 we give the *reliability* and *completeness* for selecting early- and late-type galaxies by using the *Hyperz* code (including Models A-D2) and by using the three morphology selection criteria ($C = 2.6$, $P_{\text{exp}} - P_{\text{dev}} = 0$ and $u-r = 2.22$). The definitions of *reliability* and *completeness* are from Strateva et al. (2001), i.e., the *reliability* of classification is the fraction of galaxies from the selected subsample that are correctly classified, and the *completeness* is the fraction of all galaxies of a given type from the original sample that are selected by the classification scheme.

From Tables. 4 and 5 we see that the *reliability* for early-type selection by the *Hyperz* code (including all Models) is similar to that by the three morphology selection criteria, while the *completeness* is relatively low. And, the *reliability* and *completeness* for selecting late-type galaxies (except Models C and C2) by the *Hyperz* code is less than those of the three morphology criteria.

5.4 New morphology selection criteria

Plotting the galaxies in the morphological sample in colour-colour diagrams (Fig. 27), we find that $F_{\text{UV}}-u$ and $N_{\text{UV}}-u$ can be used as new morphology indicators. In Table 6 we present the *reliability* and *completeness* for selecting early- and late-type galaxies by the $F_{\text{UV}}-u = 3.16$ and $N_{\text{UV}}-u = 1.94$ separators. We find that the $N_{\text{UV}}-u = 1.94$ colour separator has comparable *reliability* and *completeness* for both early- and late-type selection to the concentration index criterion ($C = 2.6$), and has higher *completeness* ($\sim 73\%$) for late-type selection than the $u-r = 2.22$ colour separator ($\sim 40\%$). Moreover, the $F_{\text{UV}}-u = 3.16$ colour separator has less *completeness* for selecting early-type galaxies and less *reliability* for selecting late-type galaxies than $C = 2.6$, profile likelihood $P_{\text{dev}} = P_{\text{exp}}$ and $u-r = 2.22$ separators, this is explained by that Sa-type galaxies show more scatter in $F_{\text{UV}}-u$ colour.

If the galaxies have not been detected in the N_{UV} pass-band, we can construct the morphology selection criterion by combining their optical with F_{UV} photometry, such as, we construct a morphology selection separator of $5.77-1.47(u-r) - (F_{\text{UV}}-u) = 0$. By comparison we find that this separator has comparable *reliability* and *completeness* for both early- and late-type selection to $C = 2.6$ and $N_{\text{UV}}-u = 1.94$ selection criteria (see Table. 6) and has higher *completeness* for selecting late types than $u-r = 2.22$ separator. Similarly, we also construct the $5.27-1.30(u-r) - (N_{\text{UV}}-u) = 0$ separator, and find that this criterion has similar *reliability* and *completeness* for both early- and late-type selection to $N_{\text{UV}}-u = 1.94$ criterion, i.e., the performance of early- and late-type selection has not been improved even if the optical photometry is combined with $N_{\text{UV}}-u$ colour. In Fig. 27 we present the lines, corresponding to the morphology selection separators, in the diagrams of $(F_{\text{UV}}-u)$.vs. $(u-r)$ and $(N_{\text{UV}}-u)$.vs. $(u-r)$.

6 SUMMARY AND CONCLUSIONS

Using the *Hyperz* code of Bolzonella et al. (2000) and a template spectral library which consists of 4 observed galaxy

spectra from CWW and 8 spectral families built with the GISSSEL98, BC03, HPL07 and Yunnan EPS models, respectively, we present photometric redshift estimates for a spectroscopic sample of galaxies which are selected randomly from the SDSS DR7 and GALEX DR4, and present morphological types for a morphological sample of bright galaxies which are also matched with the SDSS DR7 and GALEX DR4.

Based on the spectroscopic galaxy sample, we find that the inclusion of F_{UV} or N_{UV} or both photometry can decrease the number of catastrophic identifications ($|z_{\text{spec}} - z_{\text{phot}}| > 1.0$), and the number of catastrophic identifications decreases from in the case of using $F_{UV}N_{UV}ugriz$ to $F_{UV}ugriz$ and $N_{UV}ugriz$ photometry. In the case of using $F_{UV}N_{UV}ugriz$ photometry the catastrophic objects are those galaxies with blue $g - r$ colour and low spectroscopic redshift, in the case of using $N_{UV}ugriz$ photometry they are those faint 'redder' high-redshift galaxies, and in the case of using $F_{UV}ugriz$ photometry the catastrophic identifications are those 'bluer' low-redshift galaxies. If GALEX F_{UV} photometry or both F_{UV} and N_{UV} photometry are combined with $ugriz$ data, the difference between photometric and spectroscopic redshifts is within 0.2 for the majority of non-catastrophic objects. If only N_{UV} is combined, the derived photo- z s are systematically greater than their spectroscopic counterpart.

If catastrophic identifications are removed, the inclusion of both F_{UV} and N_{UV} photometry can raise the number of non-catastrophic identifications in the low redshift, $g - r \lesssim 0.8$ and faint r -magnitude regions, decrease the mean deviation $|\langle \Delta z \rangle|$ in the $z_{\text{spec}} \gtrsim 0.5$ and $0.3 \lesssim g - r \lesssim 0.8$ regions (except Models C and C2), and decrease the standard deviation σ_z in the $z_{\text{spec}} \gtrsim 0.3$ and $0.3 \lesssim g - r \lesssim 0.6$ regions. Moreover, if a galaxy has fainter u light, its photo- z estimate still would be erroneously identified even if UV photometry is included, and if a galaxy has fainter N_{UV} light, its photo- z estimate would be erroneously identified when UV photometry is included. Binary interactions mainly can raise the number of non-catastrophic identifications and decrease the mean and standard deviations in the $0.3 \leq g - r \leq 0.8$ region in the case of only using optical photometry.

Based on the morphological galaxy sample, we confirm that early types can be easily retrieved and late types are often misidentified as early types, and find that the inclusion of UV photometry decreases and increases the probability that early types are classified as Burst and E types, respectively, and increases the probability that late types are classified as CWW-Sbc and CWW-Scd types. If catastrophic identifications are excluded, the inclusion of UV data raises the number of the retrieved late-type galaxies in all redshift, bluer $g - r$ and $r \gtrsim 14$ regions, and the increment increases with $T(\text{Fuk})$. Binary interactions mainly affect the determination of E and S0 types.

By comparison we find the *reliability* and *completeness* for early- and late-type selection by the *Hyperz* code are much lower than those by the three morphology criteria: concentration index $C = 2.6$, profile likelihood $P_{\text{dev}} = P_{\text{exp}}$ and $u - r = 2.22$. Moreover, we find that $N_{UV} - u = 1.94$ can be used as a new morphology selection indicator, its *reliability* and *completeness* for selecting early and late types are approximately the same as those of $C = 2.6$ criterion, and even its *completeness* for selecting early types is greater than

that of $u - r = 2.22$ criterion. The criterion of $F_{UV} - u = 3.16$ is not as good as $N_{UV} - u = 1.94$, while the criterion constructed by the combination of $F_{UV} - u$ with $u - r$ colour, $5.77 - 1.47(u - r) = F_{UV} - u$, can reach to the similar *reliability* and *completeness* for early- and late-type selection to those of $N_{UV} - u = 1.94$. For the new constructed criteria $5.27 - 1.30(u - r) = N_{UV} - u$, the *reliability* and *completeness* for early- and late-type selection are not been improved in comparison with $N_{UV} - u = 1.94$ criterion.

ACKNOWLEDGMENTS

This work was funded by the Chinese Natural Science Foundation (Grant Nos 10773026, 10673029, 10821026 & 2007CB15406) and by Yunnan Natural Science Foundation (Grant No 2007A113M). We are also grateful to the referee for suggestions that have improved the quality of this manuscript.

Funding for the SETS and SDSS-II has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, the U.S. Department of Energy, the National Aeronautics and Space Administration, the Japanese Monbukagakusho, the Max Planck Society, and the Higher Education Funding Council for England. The SDSS Web Site is <http://www.sdss.org/>.

The SDSS is managed by the Astrophysical Research Consortium for the Participating Institutions. The Participating Institutions are the American Museum of Natural History, Astrophysical Institute Potsdam, University of Basel, University of Cambridge, Case Western Reserve University, University of Chicago, Drexel University, Fermilab, the Institute for Advanced Study, the Japan Participation Group, Johns Hopkins University, the Joint Institute for Nuclear Astrophysics, the Kavli Institute for Particle Astrophysics and Cosmology, the Korean Scientist Group, the Chinese Academy of Sciences (LAMOST), Los Alamos National Laboratory, the Max-Planck-Institute for Astronomy (MPIA), the Max-Planck-Institute for Astrophysics (MPA), New Mexico State University, Ohio State University, University of Pittsburgh, University of Portsmouth, Princeton University, the United States Naval Observatory, and the University of Washington.

The galaxy Evolution Explorer (GALEX) is a NASA Small Explorer. The mission was developed in cooperation with the centre National d'Etudes Spatiales of France and the Korean Ministry of Science and Technology.

REFERENCES

- Abdalla F., Banerji M., Lahav O., Rashkov V., 2008, arXiv0812.3831
- Agueros M. A., Ivezić A., Covey K. R., Obric M., Hao L., Walkowicz L. M., West A. A., Vanden Berk D. E., et al., 2005, ApJ, 130, 1022
- Assef R., Kochanek C., Brodwin M., Brown M., Caldwell N., Cool R., Eisenhardt P., Eisenstein D., et al., 2008, ApJ, 676, 286
- Babbedge T., Rowan-Robinson M., Gonzalez-Solares E., Polletta M., Berta S., Perez-Fournon I., Oliver S., Salaman D., et al., 2004, MNRAS, 353, 654

- Baum, W., 1962, in George C. M., eds, IAU Symp. 15, Problems of Extra-Galactic Research. Macmillan Press, New York, p.390
- Benitez N., 2000, ApJ, 536, 571
- Blanton M., Dalcanton J., Eisenstein D., Loveday J., Strauss M., SubbaRao M., Weinberg D., Anderson J., et al. 2001, AJ, 121, 2358
- Blanton M., Roweis S., 2007, AJ, 133, 734
- Bolzonella M., Miralles J. M., Pelló R., 2000, A&A, 363, 476
- Brammer G., van Dokkum P., Coppi P., 2008, ApJ, 686, 1053
- Bruzual G., Charlot S., 1993, ApJ, 405, 538
- Bruzual G., Charlot S., 2003, MNRAS, 344, 1000 [BC03]
- Calzetti D., Armus L., Bohlin R. C., Kinney A. L., Koornneef J., Storchi-Bergmann T., 2000, ApJ, 533, 682
- Cardelli J. A., Clayton G., Mathis J. S., 1989, ApJ, 345, 245 [CCM]
- Chabrier G., 2003, PASP, 115, 763
- Coleman G., Wu C., Weedman D., 1980, ApJS, 43, 393
- Donas J., Deharveng J., Rich R. M., Yi, S. K., Lee Y., Boselli A., Gil de Paz A., Boissier S., et al., 2007, ApJS, 173, 597
- Feldmann R., Carollo C., Porciani C., Lilly S., Capak P., Taniguchi Y., Le Fevre O., Renzini A., et al., 2006, MNRAS, 372, 565
- Fitzpatrick E. L., 1986, AJ, 92, 1068
- Fukugita M., Nakamura O., Okamura S., Yasuda N., Barntine J., Brinkmann J., Gunn J., Harvanek M., et al. 2007, AJ, 134, 579
- Gerdes D., Sypniewski A., McKay T., Hao J., Weis M., Wechsler R., Busha M., 2009, arXiv0908.4085
- Han Z., Podsiadlowski Ph., Maxted P. F. L., Marsh T. R., Ivanova N., 2002, MNRAS, 336, 449
- Han Z., Podsiadlowski Ph., Maxted P. F. L., Marsh T. R., 2003, MNRAS, 341, 669
- Han Z., Podsiadlowski Ph., Lynas-Gray A. E., 2007, MNRAS, 380, 1098 [HPL07]
- Hurley J. R., Pols O. R., Tout C. A., 2000, MNRAS, 315, 543
- Hurley J. R., Tout C. A., Pols O. R., 2002, MNRAS, 329, 897
- Kurtz M. J., Geller M. J., Fabricant D. G., Wyatt W. F., Dell Antonio I. P., 2007, AJ, 134, 1360
- Lilly S. J., Le Fevre O., Renzini A., Zamorani G., Scodeggio M., Contini T., Carollo C. M., Hasinger G., et al., 2007, ApJS, 172, 70
- Lintott C., Schawinski K., Slosar A., Land K., Bamford S., Thomas D., Raddick M., Nichol R., et al., 2008, MNRAS, 389, 1179
- Lupton R. H., 2001, ASPC, 232, 2001
- Maraston C., 2005, MNRAS, 362, 799
- Miller G. E., Scalo J. M., 1979, ApJS, 41, 513
- Mobasher B., Capak P., Scoville N., Dahlen T., Salvato M., Aussel H., Thompson D., Feldmann R., et al., 2007, ApJS, 172, 117
- Negrello M., Serjeant S., Pearson C., Takagi T., Efstathiou A., Goto, T., Burgarella D., et al. 2009, MNRAS, 394, 375
- Niemack M., Jimenez R., Verde L., Menanteau F., Panter B., Spergel D., 2009, ApJ, 690, 89
- Obric M., Ivezić Z., Best P. N., Lupton R. H., Tremonti C., Brinchmann J., Agueros M. A., Knapp G. R., et al., 2006, MNRAS, 370, 1677
- Rich R. M., Salim S., Brinchmann J., Charlot S., Seibert M., Kauffmann G., Lee Y. et al. 2005, ApJ, 619, 107
- Rudnick G., Franx M., Rix H., Moorwood A., Kuijken K., van Starkenburg L., van der Werf P., Rottgering H., et al. 2001, AJ, 122, 2205
- Rudnick G., Rix H., Franx M., Labbe I., Blanton M., Daddi E., Forster Schreiber N., Moorwood A., et al. 2003, ApJ, 599, 847
- Sandage A., 1961, The Hubble Atlas of Galaxies (Washington: Carnegie Inst. Washington)
- Sarajedini V. L., Green R. F., Griffiths R. E., Ratnatunga K. 1999, ApJS, 121, 417
- Seaton M. J., 1979, MNRAS, 187, 73
- Shimasaku K., Fukugita M., Doi M., Hamabe M., Ichikawa T., Okamura S., Sekiguchi M., Yasuda N., Brinkmann J., 2001, AJ, 122, 1238
- Schlegel D. J., Finkbeiner D., Douglas P., Davis M., 1998, ApJ, 500, 525
- Schmidt M., 1959, ApJS, 129, 243
- Strateva I., Ivezić Z., Knapp G., Narayanan V. K., Strauss M. A., Gunn J. E., Lupton R. H., Schlegel D., et al., 2001, AJ, 122, 1861
- Wang T., Huang, J., Gu, Q., 2009, RAA, 9, 390
- Wray J. J. & Gunn J. E., 2008, ApJ, 678, 144
- Wuyts S., Franx M., Cox T., Hernquist L., Hopkins Ph.,; Robertson B., van Dokkum P., 2009, ApJ, 696, 348
- Wyder T., Treyer M., Milliard B., Schiminovich D., Arnouts S., Budavari T., Barlow T., Bianchi L., et al. 2005, ApJ, 619, 15
- Xia L., Cohen S., Malhotra S., Rhoads J., Grogin N., Hathi N., Windhorst R., et al. 2009, AJ, 138, 95
- York D. G., Adelman J., Anderson J. E. Jr., Anderson S. F., Annis J., Bahcall N. A., Bakken J. A., Barkhouser R., et al. 2000, AJ, 120, 1579
- Zhang F., Han, Z., Li, L., Hurley J. R., 2002, MNRAS, 334, 883
- Zhang F., Han, Z., Li, L., Hurley J. R., 2004a, MNRAS, 350, 710
- Zhang F., Han, Z., Li, L., Hurley J. R., 2004b, A&A, 415, 117
- Zhang F., Han, Z., Li, L., Hurley J. R., 2005, MNRAS, 357, 1088
- Zhang Y., Li L., Zhao Y., 2009, MNRAS, 392, 233

APPENDIX A: THE INFLUENCE OF INPUT PARAMETERS ON PHOTO-Z DETERMINATIONS.

In this section our standard is Model B. To investigate the effect of the input parameters on photo-z estimates we proceed by changing each parameter, in turn, while the other parameters remain the same as in the standard model. Model Ba uses the reddening law of Seaton (1979) instead of Calzetti et al. (2000). Model Bb uses the set of minimum magnitudes [26, 26, 24, 24, 24, 24, 24] instead of [29.5, 29.5, 30, 30, 30, 30, 30]. Models Bc and Bd use the minimum magnitude error of 0.001 and 0.10 instead of 0.05. Models Be and Bf only use the 8 theoretical spectral families and CWW spectra, respectively. Model Bg uses the set of cosmological parameters [1.0, 0.0, 50] instead of [0.3, 0.7, 70].




Figure A1. The differences in N (top panels), $|\langle\Delta z\rangle|$ (middle panels) and σ_z (bottom panels) between Model B and the other models Ba-Bg as a function of z_{spec} in removing catastrophic identifications. Left and right panels are for using *ugriz* and $F_{\text{UV}}N_{\text{UV}}\textit{ugriz}$ photometry, respectively.

We obtain photo- z estimates of galaxies in the spectroscopic sample for Models Ba-Bf by using *ugriz* and $F_{\text{UV}}N_{\text{UV}}\textit{ugriz}$ photometry. In Fig. A1-A3 we present the differences in the number of non-catastrophic identifications (ΔN), mean ($\Delta|\langle\Delta z\rangle|$) and standard deviations ($\Delta\sigma_z$) between Model B and the other models Ba-Bg as a function of redshift, $g-r$ colour and r -magnitude in removing catastrophic identifications, respectively. From them we see that

- if the reddening law of Seaton (1979) [Model Ba] is used, the number of non-catastrophic identifications N increases in the case of using optical photometry, the mean and standard deviations (except in the case of using $F_{\text{UV}}N_{\text{UV}}\textit{ugriz}$ photometry) significantly differ from those of Model B;
- if the minimum magnitude error is set to 0.001 (Model Bc, one-fifty of the value in Model B) the number of non-catastrophic identifications N , the mean and standard deviations in the bluer $g-r$ region are greater than those of Model B in the case of only using optical photometry;
- if only theoretical spectral template (Model Be) or only the CWW set of empirical spectra (Model Bf) is used, the number of non-catastrophic identifications would be increased or decreased in the case of using optical photometry, respectively, the standard deviation is not changed significantly, and the mean deviation of Model Bf is greater than that of Model Be;
- decreasing the set of minimum magnitude (Model Bb) or increasing the minimum magnitude error by a factor of 2 (0.1, Model Bd) or using the set of cosmological parameters of [1.0, 0., 70] would not vary significantly the results.




Figure A2. Similar to Fig. A1, but a function of $g-r$ colour.




Figure A3. Similar to Fig. A1, but a function of r -magnitude.

APPENDIX B: DEPENDENCE OF PHOTO- z ACCURACY ON REDSHIFT, COLOUR, MAGNITUDE AND THE FILTER SET.

In Tables B1-B3 we give the number of non-catastrophic identifications N , standard deviation σ_z and mean deviation $\langle\Delta z\rangle$ as a function of spectroscopic redshift z_{spec} , $g-r$ colour and r -magnitude in removing catastrophic identifications for Models A-D2. Each table includes the cases of using *ugriz*, $F_{\text{UV}}N_{\text{UV}}\textit{ugriz}$, $F_{\text{UV}}\textit{ugriz}$ (only for Model B) and $N_{\text{UV}}\textit{ugriz}$ (only for Model B) photometry.

APPENDIX C: DEPENDENCE OF MORPHOLOGY ACCURACY ON REDSHIFT, COLOUR AND MAGNITUDE.

In Figs.C1-C3 we illustrate the evolutions of N_i , $|\langle\Delta T_i\rangle|$ and σ_{T_i} as a function of spectroscopic redshift z_{spec} , $g-r$ colour and r -magnitude in removing catastrophic identifications for

Table B1. The number of non-catastrophic identifications N , standard deviation (σ_z) and mean deviation ($\langle\Delta z\rangle$) of galaxies in the spectroscopic sample as a function of the redshift bin and the filter set for Models A-D2. The first line gives the number of galaxies in the corresponding redshift bin. The following 1st and 2nd parts give the results when $ugriz$ and $F_{UV}N_{UV}ugriz$ photometry is used for all models, and the 3rd and 4th parts give the results when $N_{UV}ugriz$ and $F_{UV}ugriz$ photometry is used for Model B. All results are in the cases of removing catastrophic identifications.

	$z = 0.0 - 0.05$ $N, \sigma_z, \langle\Delta z\rangle$	$z = 0.05 - 0.10$ $N, \sigma_z, \langle\Delta z\rangle$	$z = 0.10 - 0.15$ $N, \sigma_z, \langle\Delta z\rangle$	$z = 0.15 - 0.20$ $N, \sigma_z, \langle\Delta z\rangle$	$z = 0.20 - 0.30$ $N, \sigma_z, \langle\Delta z\rangle$	$z = 0.30 - 0.40$ $N, \sigma_z, \langle\Delta z\rangle$	$z = 0.40 - 0.50$ $N, \sigma_z, \langle\Delta z\rangle$	$z = 0.50 - 0.60$ $N, \sigma_z, \langle\Delta z\rangle$
All	1114	2395	1796	783	288	101	47	5
				$ugriz, non-cata.$				
A	722, 0.054,-0.035	1658, 0.041, 0.003	1573, 0.045, 0.017	765, 0.045, 0.011	279, 0.056, 0.004	87, 0.085,-0.047	41, 0.082, 0.017	5, 0.064,-0.032
B	741, 0.054,-0.022	1881, 0.040, 0.012	1680, 0.045, 0.025	771, 0.046, 0.014	279, 0.064,-0.007	87, 0.077,-0.061	41, 0.086,-0.001	5, 0.066,-0.022
C	673, 0.061,-0.060	1471, 0.058,-0.022	1407, 0.060,-0.022	724, 0.055,-0.027	273, 0.062,-0.036	86, 0.069,-0.058	40, 0.086,-0.016	5, 0.067,-0.019
C2	666, 0.061,-0.060	1464, 0.059,-0.021	1405, 0.061,-0.020	723, 0.056,-0.025	271, 0.061,-0.038	80, 0.069,-0.065	40, 0.086,-0.016	4, 0.058,-0.047
D	693, 0.058,-0.042	1567, 0.044, 0.001	1538, 0.049, 0.012	756, 0.048, 0.008	278, 0.058, 0.002	87, 0.085,-0.050	41, 0.092, 0.015	5, 0.104, 0.003
D2	706, 0.055,-0.039	1609, 0.042, 0.003	1550, 0.046, 0.015	738, 0.046, 0.011	250, 0.058,-0.005	81, 0.082,-0.052	35, 0.086, 0.016	4, 0.051,-0.050
				$F_{UV}N_{UV}ugriz, non-cata.$				
A	1109, 0.056,-0.027	2385, 0.049, 0.009	1790, 0.052, 0.021	780, 0.049, 0.016	283, 0.059,-0.001	94, 0.073,-0.040	44, 0.050, 0.015	5, 0.080, 0.019
B	1109, 0.055,-0.023	2384, 0.049, 0.016	1790, 0.055, 0.027	781, 0.056, 0.014	283, 0.064,-0.008	93, 0.081,-0.050	44, 0.049, 0.018	5, 0.081, 0.021
C	1111, 0.066,-0.043	2382, 0.064,-0.021	1791, 0.066,-0.027	780, 0.064,-0.035	283, 0.069,-0.051	92, 0.071,-0.070	42, 0.066,-0.040	5, 0.067,-0.053
C2	1110, 0.068,-0.045	2378, 0.065,-0.020	1791, 0.068,-0.025	779, 0.065,-0.033	283, 0.070,-0.052	90, 0.069,-0.069	42, 0.066,-0.032	4, 0.063,-0.070
D	1109, 0.061,-0.038	2383, 0.053, 0.001	1790, 0.053, 0.013	780, 0.051, 0.014	283, 0.058,-0.003	93, 0.079,-0.044	44, 0.048, 0.020	5, 0.103, 0.038
D2	1109, 0.061,-0.036	2381, 0.052, 0.003	1788, 0.052, 0.015	777, 0.051, 0.012	279, 0.061,-0.007	91, 0.078,-0.052	41, 0.050, 0.018	4, 0.084, 0.000
				$N_{UV}ugriz, non-cata.$				
B	1091, 0.107,-0.367	2337, 0.101,-0.361	1779, 0.100,-0.400	765, 0.097,-0.434	236, 0.088,-0.450	42, 0.082,-0.440	14, 0.123,-0.457	1, ∞ ,-0.337
				$F_{UV}ugriz, non-cata.$				
B	1090, 0.063,-0.033	2334, 0.046, 0.013	1781, 0.051, 0.020	776, 0.049, 0.010	280, 0.058,-0.011	88, 0.083,-0.051	42, 0.076, 0.002	5, 0.066,-0.022

Table B2. Similar to Table B1, but for the bin of $g-r$ colour.


	$g-r=0.0-0.2$ $N, \sigma_z, \langle\Delta z\rangle$	$g-r=0.2-0.4$ $N, \sigma_z, \langle\Delta z\rangle$	$g-r=0.4-0.6$ $N, \sigma_z, \langle\Delta z\rangle$	$g-r=0.6-0.8$ $N, \sigma_z, \langle\Delta z\rangle$	$g-r=0.8-1.0$ $N, \sigma_z, \langle\Delta z\rangle$	$g-r=1.0-1.2$ $N, \sigma_z, \langle\Delta z\rangle$	$g-r=1.2-1.4$ $N, \sigma_z, \langle\Delta z\rangle$	$g-r=1.4-1.6$ $N, \sigma_z, \langle\Delta z\rangle$
All	17	344	1109	1628	2050	897	230	91 (160)
				$ugriz, non-cata.$				
A	1, ∞ ,-0.003	64, 0.059,-0.041	486, 0.067,-0.028	1212, 0.044,-0.011	2018, 0.040, 0.015	896, 0.046, 0.018	228, 0.044, 0.008	86, 0.081,-0.023
B	3, 0.004, 0.007	74, 0.041, 0.001	680, 0.053,-0.010	1351, 0.045, 0.006	2027, 0.040, 0.024	896, 0.048, 0.014	228, 0.052,-0.006	87, 0.081,-0.051
C	8, 0.030,-0.022	94, 0.072,-0.048	374, 0.075,-0.031	836, 0.055,-0.037	2025, 0.057,-0.030	895, 0.062,-0.016	228, 0.064,-0.020	86, 0.083,-0.047
C2	10, 0.049,-0.036	86, 0.078,-0.051	367, 0.076,-0.030	835, 0.055,-0.036	2023, 0.057,-0.029	894, 0.063,-0.013	228, 0.064,-0.018	83, 0.079,-0.060
D	1, ∞ ,-0.055	38, 0.069,-0.066	398, 0.073,-0.037	1160, 0.046,-0.020	2023, 0.042, 0.012	896, 0.049, 0.018	228, 0.053, 0.010	84, 0.082,-0.019
D2	2, 0.003,-0.051	51, 0.069,-0.040	429, 0.071,-0.029	1195, 0.044,-0.013	2018, 0.041, 0.013	891, 0.049, 0.017	197, 0.047, 0.008	70, 0.090,-0.050
				$F_{UV}N_{UV}ugriz, non-cata.$				
A	16, 0.050,-0.045	344, 0.053,-0.023	1104, 0.060,-0.013	1622, 0.054, 0.007	2041, 0.048, 0.019	897, 0.048, 0.020	228, 0.052, 0.000	90, 0.077,-0.036
B	16, 0.059,-0.044	344, 0.053,-0.005	1105, 0.062, 0.002	1621, 0.056, 0.014	2040, 0.052, 0.021	897, 0.052, 0.015	228, 0.056,-0.010	91, 0.081,-0.045
C	17, 0.048,-0.037	344, 0.068,-0.040	1105, 0.067,-0.033	1621, 0.069,-0.031	2040, 0.063,-0.023	897, 0.062,-0.025	228, 0.074,-0.037	89, 0.080,-0.068
C2	17, 0.054,-0.043	344, 0.068,-0.040	1105, 0.069,-0.033	1620, 0.069,-0.031	2035, 0.064,-0.022	897, 0.064,-0.021	228, 0.074,-0.034	89, 0.080,-0.071
D	16, 0.053,-0.091	344, 0.056,-0.046	1105, 0.061,-0.024	1621, 0.055,-0.003	2039, 0.050, 0.014	897, 0.047, 0.020	228, 0.060, 0.004	90, 0.092,-0.031
D2	16, 0.054,-0.087	344, 0.056,-0.044	1105, 0.062,-0.021	1621, 0.055, 0.001	2037, 0.049, 0.014	894, 0.047, 0.018	224, 0.061, 0.001	89, 0.087,-0.047
				$N_{UV}ugriz, non-cata.$				
B	17, 0.084,-0.392	342, 0.086,-0.359	1105, 0.100,-0.321	1619, 0.109,-0.350	1980, 0.079,-0.410	880, 0.080,-0.451	207, 0.071,-0.506	61, 0.112,-0.485
				$F_{UV}ugriz, non-cata.$				
B	17, 0.102,-0.070	339, 0.056,-0.014	1094, 0.065,-0.011	1563, 0.056, 0.004	2032, 0.045, 0.020	896, 0.049, 0.013	228, 0.048,-0.006	87, 0.073,-0.046

Models A-D2. This part is based on morphological galaxy sample.

This paper has been typeset from a $\text{\LaTeX}$ file prepared by the author.

Table B3. Similar to Table B1, but for the r -magnitude bin.

	$r=15.5-16.0$ $N, \sigma_z, \langle \Delta z \rangle$	$r=16.0-16.5$ $N, \sigma_z, \langle \Delta z \rangle$	$r=16.5-17.0$ $N, \sigma_z, \langle \Delta z \rangle$	$r=17.0-17.5$ $N, \sigma_z, \langle \Delta z \rangle$	$r=17.5-18.0$ $N, \sigma_z, \langle \Delta z \rangle$	$r=18.0-18.5$ $N, \sigma_z, \langle \Delta z \rangle$	$r=18.5-19.0$ $N, \sigma_z, \langle \Delta z \rangle$	$r=19.0-19.5$ $N, \sigma_z, \langle \Delta z \rangle$
All	(130) 149	275	478	929	1498	1952	834	96 (187)
	<i>ugriz</i> , non-cata.							
A	117, 0.041,-0.005	212, 0.051,-0.008	358, 0.043, 0.003	660, 0.047,-0.002	1062, 0.050, 0.002	1536, 0.050, 0.008	825, 0.046, 0.010	94, 0.079,-0.011
B	124, 0.041, 0.001	225, 0.045, 0.005	383, 0.036, 0.014	726, 0.046, 0.009	1171, 0.049, 0.012	1674, 0.046, 0.016	825, 0.049, 0.011	94, 0.077,-0.028
C	122, 0.061,-0.039	202, 0.061,-0.027	328, 0.057,-0.019	583, 0.060,-0.028	945, 0.062,-0.029	1345, 0.060,-0.027	809, 0.059,-0.031	94, 0.077,-0.037
C2	120, 0.059,-0.039	201, 0.062,-0.029	326, 0.058,-0.018	579, 0.061,-0.025	944, 0.062,-0.028	1340, 0.061,-0.026	809, 0.060,-0.030	92, 0.076,-0.038
D	115, 0.048,-0.010	208, 0.054,-0.008	351, 0.048,-0.001	627, 0.051,-0.003	1013, 0.054,-0.002	1478, 0.053, 0.003	819, 0.049, 0.008	94, 0.085,-0.007
D2	118, 0.047,-0.010	213, 0.051,-0.013	355, 0.043, 0.001	638, 0.049,-0.001	1035, 0.051, 0.002	1491, 0.049, 0.007	803, 0.049, 0.008	78, 0.083,-0.022
	$F_{UV} N_{UV} \text{ugriz}$, non-cata.							
A	149, 0.052,-0.015	274, 0.054,-0.009	478, 0.056, 0.001	927, 0.049, 0.004	1489, 0.054, 0.006	1944, 0.053, 0.015	831, 0.049, 0.016	96, 0.067,-0.018
B	149, 0.054,-0.015	273, 0.053,-0.005	478, 0.056, 0.004	927, 0.049, 0.011	1489, 0.054, 0.014	1945, 0.057, 0.021	831, 0.053, 0.014	96, 0.075,-0.025
C	149, 0.069,-0.040	273, 0.070,-0.029	478, 0.065,-0.021	927, 0.065,-0.027	1490, 0.065,-0.028	1944, 0.064,-0.024	831, 0.062,-0.037	96, 0.086,-0.060
C2	149, 0.074,-0.039	270, 0.072,-0.032	476, 0.067,-0.020	927, 0.066,-0.027	1490, 0.066,-0.027	1944, 0.065,-0.023	831, 0.063,-0.035	96, 0.086,-0.061
D	149, 0.060,-0.023	272, 0.059,-0.017	477, 0.060,-0.007	928, 0.053,-0.004	1490, 0.057,-0.002	1944, 0.055, 0.007	831, 0.048, 0.013	96, 0.081,-0.018
D2	149, 0.057,-0.023	271, 0.056,-0.014	476, 0.059,-0.006	928, 0.054,-0.003	1488, 0.056, 0.000	1944, 0.054, 0.009	827, 0.050, 0.013	95, 0.080,-0.028
	$N_{UV} \text{ugriz}$, non-cata.							
B	137, 0.097,-0.381	258, 0.099,-0.374	456, 0.090,-0.367	923, 0.096,-0.366	1482, 0.102,-0.375	1927, 0.106,-0.378	814, 0.095,-0.441	73, 0.083,-0.516
	$F_{UV} \text{ugriz}$, non-cata.							
B	143, 0.052,-0.006	268, 0.052, 0.000	471, 0.057, 0.005	918, 0.054, 0.004	1472, 0.055, 0.007	1913, 0.055, 0.010	830, 0.051, 0.011	94, 0.070,-0.026

**Figure C1.** Evolutions of N_i (green dotted line), $|\Delta T_i|$ (black solid lines) and σ_{T_i} (red dashed lines) as a function of spectroscopic redshift z_{spec} in both cases of using *ugriz* (thin lines) and $F_{UV} N_{UV} \text{ugriz}$ (thick lines) photometry. In the shaded regions $N_i \geq 10$.




Figure C2. Similar to Fig. C1, but as a function of $g-r$ colour.




Figure C3. Similar to Fig. C1, but as a function of r -magnitude.

